

JC20 Rec'd PCT/PTO 13 MAR 2002

ORIGINAL

FORM PTO- 390 (REV -2000)		U.S. DEPARTMENT OF COMMERCE PATENT AND TRADEMARK OFFICE		ATTORNEY'S DOCKET NUMBER Le A 33 965	
TRANSMITTAL LETTER TO THE UNITED STATES DESIGNATED/ELECTED OFFICE (DO/EO/US) CONCERNING A FILING UNDER 35 U.S.C. 371				U.S. APPLICATION NO. (If known, see 37 CFR 1.5)	
				10/070963	
INTERNATIONAL APPLICATION NO. PCT/EP00/08836		INTERNATIONAL FILING DATE 11 September 2000 (11.09.00)		PRIORITY DATE CLAIMED 15 September 1999 (15.09.99)	
TITLE OF INVENTION NOVEL COMBINATION FOR THE TREATMENT OF SEXUAL DYSFUNCTION					
APPLICANT(S) FOR DO/EO/US BISCHOFF, Erwin; BISCHOFF, Hilmar; GIULIANO, Francois					
Applicant herewith submits to the United States Designated/Elected Office (DO/EO/US) the following items and other information:					
<p>1 <input checked="" type="checkbox"/> This is a FIRST submission of items concerning a filing under 35 U.S.C. 371.</p> <p>2 <input type="checkbox"/> This is a SECOND or SUBSEQUENT submission of items concerning a filing under 35 U.S.C. 371.</p> <p>3 <input checked="" type="checkbox"/> This is an express request to begin national examination procedures (35 U.S.C. 371(f)). The submission must include items (5), (6), (9) and (21) indicated below.</p> <p>4 <input checked="" type="checkbox"/> The US has been elected by the expiration of 19 months from the priority date (Article 31).</p> <p>5 <input checked="" type="checkbox"/> A copy of the International Application as filed (35 U.S.C. 371(c)(2))</p> <p style="margin-left: 20px;">a. <input checked="" type="checkbox"/> is attached hereto (required only if not communicated by the International Bureau).</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> has been communicated by the International Bureau.</p> <p style="margin-left: 20px;">c. <input type="checkbox"/> is not required, as the application was filed in the United States Receiving Office (RO/US).</p> <p>6 <input checked="" type="checkbox"/> An English language translation of the International Application as filed (35 U.S.C. 371(c)(2)).</p> <p style="margin-left: 20px;">a. <input checked="" type="checkbox"/> is attached hereto.</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> has been previously submitted under 35 U.S.C. 154(d)(4).</p> <p>7 <input checked="" type="checkbox"/> Amendments to the claims of the International Application under PCT Article 19 (35 U.S.C. 371(c)(3))</p> <p style="margin-left: 20px;">a. <input type="checkbox"/> are attached hereto (required only if not communicated by the International Bureau).</p> <p style="margin-left: 20px;">b. <input type="checkbox"/> have been communicated by the International Bureau.</p> <p style="margin-left: 20px;">c. <input type="checkbox"/> have not been made; however, the time limit for making such amendments has NOT expired.</p> <p style="margin-left: 20px;">d. <input checked="" type="checkbox"/> have not been made and will not be made.</p> <p>8 <input type="checkbox"/> An English language translation of the amendments to the claims under PCT Article 19 (35 U.S.C. 371 (c)(3)).</p> <p>9 <input type="checkbox"/> An oath or declaration of the inventor(s) (35 U.S.C. 371(c)(4)).</p> <p>10 <input type="checkbox"/> An English language translation of the annexes of the International Preliminary Examination Report under PCT Article 36 (35 U.S.C. 371(c)(5)).</p> <p>Items 11 to 20 below concern document(s) or information included:</p> <p>11 <input checked="" type="checkbox"/> An Information Disclosure Statement under 37 CFR 1.97 and 1.98.</p> <p>12 <input type="checkbox"/> An assignment document for recording. A separate cover sheet in compliance with 37 CFR 3.28 and 3.31 is included.</p> <p>13 <input checked="" type="checkbox"/> A FIRST preliminary amendment.</p> <p>14 <input type="checkbox"/> A SECOND or SUBSEQUENT preliminary amendment.</p> <p>15 <input type="checkbox"/> A substitute specification.</p> <p>16 <input type="checkbox"/> A change of power of attorney and/or address letter.</p> <p>17 <input type="checkbox"/> A computer-readable form of the sequence listing in accordance with PCT Rule 13ter.2 and 35 U.S.C. 1.821 - 1.825.</p> <p>18 <input type="checkbox"/> A second copy of the published international application under 35 U.S.C. 154(d)(4).</p> <p>19 <input type="checkbox"/> A second copy of the English language translation of the international application under 35 U.S.C. 154(d)(4).</p> <p>20 <input checked="" type="checkbox"/> Other items or information: 1) Certificate of Mailing under 37 C.F.R. 1.10; 2) Transmittal of Information Disclosure Statement under 37 C.F.R. 1.97(b); 3) Information Disclosure Citation (Modified Form PTO-1449) and references cited therein; 4) Return Receipt Postcard.</p>					
Date of Deposit: MAR 13 2002					
Express Mail Label No.: ET131980209US					

FORM PTO-1390 (REV 11-2000) page 2 of 2

10/070963
JC13 Rec'd PCT/PTO 13 MAR 2002

Atty. Docket No.: Le A 33 965

IN THE UNITED STATES PATENT AND TRADEMARK OFFICE

APPLICANTS: Bischoff, et al.

SERIAL NO.: National Stage filing for PCT/EP00/08836

FILING DATE: herewith

TITLE: Novel Combination for the Treatment of Sexual Dysfunction

PRELIMINARY AMENDMENT

Assistant Commissioner for Patents
Washington, D.C. 20231

Sir:

This Preliminary Amendment is submitted in the above-captioned application. Please amend the application as follows:

In the Claims

Please cancel claims 2, 3, and 4.

Please amend claims 1, 5-8, and 11-21 as shown in the attached sheets. A marked version of the claim set showing the changes made is also attached.

Please add new claims 22-26 as shown in the attached sheet.

Remarks

By way of this Preliminary Amendment, claims 1-26 are pending. Claims 1, 5-8, and 11-21 have been amended. Claims 2, 3, and 4 have been cancelled. New claims 22-26 have been added. These claim amendments and additions are being made solely for purposes of

placing the claims in a format appropriate for U.S. prosecution. Applicants submit that the amendments do not change the scope of the claims as originally filed. Such amendments are therefore made to address formalities in the claim format and are not related to the patentability of the subject matter of the claims. No new matter was added by way of these claim amendments and additions.

Conclusion

Applicants believe that the subject matter of the pending claims is patentable and that the instant application should accordingly be allowed. If the Examiner believes that a conversation with Applicants' attorney would be helpful in expediting prosecution of this application, the Examiner is invited to call the undersigned attorney at (203) 812-6450.

Respectfully submitted,

Dated: March 12, 2002


Bayer Corporation
400 Morgan Lane
West Haven, CT 06516
(Tel) (203) 812-6450
(Fax) (203) 812-6459
e-mail: susan.pellegrino.b@bayer.com

Susan M. Pellegrino
Susan M. Pellegrino
Attorney for Applicants
Reg. No. 48,972

Amended Claims (Attorney Docket No. Le A 33 965)

1. A combination preparation, comprising as pharmaceutically active ingredients at least one active compound component A and at least one active compound component B, characterized in that the active compound component A is a PDE inhibitor, and the active compound component B is an antilipemic.
2. Cancelled.
3. Cancelled.
4. Cancelled.
5. (Amended) The combination preparation as claimed in claim 1, characterized in that the active compound components A and B are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
6. (Amended) The combination preparation as claimed in claim 1, characterized in that the active compound components A and B are (spatially) separated, in particular as a kit-of-parts.
7. (Amended) The combination preparation as claimed in claim 1, characterized in that the antilipemic (active compound component B) is selected from the group consisting of (a) HMG-CoA-reductase inhibitors; (b) squalene synthase inhibitors; (c) bile acid sequestrants; (d) fibric acid and its derivatives; (e) nicotinic acid and its analogs; (f) ω 3-fatty acids.
8. (Amended) The combination preparation as claimed in claim 7, characterized in that the antilipemic (active compound component B) is an HMG-CoA-reductase inhibitor.

11. (Amended) The combination preparation as claimed in claim 1, characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and is selected from the group consisting of pyrazolopyrimidones of the general formula below



in which

- R^1 represents hydrogen; C_1 - C_3 -alkyl; C_1 - C_3 -perfluoroalkyl; or C_3 - C_5 -cycloalkyl;
 R^2 denotes hydrogen; C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl; C_1 - C_3 -perfluoroalkyl; or C_3 - C_6 -cycloalkyl;
 R^3 is C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl; C_1 - C_6 -perfluoroalkyl, C_3 - C_5 -cycloalkyl; C_3 - C_6 -alkenyl; or C_3 - C_6 -alkinyl;
 R^4 represents C_1 - C_4 -alkyl, optionally substituted by OH, NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkenyl, optionally substituted by CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkanoyl, optionally substituted by NR^5R^6 ; (hydroxy)- C_2 - C_4 -alkyl, optionally substituted by NR^5R^6 , (C_2 - C_3 -alkoxy)- C_1 - C_2 -alkyl, optionally substituted by OH or NR^5R^6 , CO_2R^7 ; halogen; NR^5R^6 , $NHSO_2NR^5R^6$; $NHSO_2R^8$; $SO_2NR^9R^{10}$; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, each of which is optionally substituted by methyl;

R^5 and R^6 each independently of one another denote hydrogen or C_1 - C_4 -alkyl; or together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4- $N(R^{11})$ -piperazinyl or imidazolyl group, where this group is optionally substituted by methyl or OH;

R^7 is hydrogen or C_1 - C_4 -alkyl;

R^8 represents C_1 - C_3 -alkyl, optionally substituted by NR^5R^6 ;

R^9 and R^{10} together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4- $N(R^{12})$ -piperazinyl group, where this group is optionally substituted by C_1 - C_4 -alkyl, C_1 - C_3 -alkoxy, $NR^{13}R^{14}$ or $CONR^{13}R^{14}$;


R^{11} denotes hydrogen, C_1 - C_3 -alkyl, optionally substituted by phenyl; (hydroxy)- C_2 - C_3 -alkyl; or C_1 - C_4 -alkanoyl;

R^{12} is hydrogen, C_1 - C_6 -alkyl, (C_1 - C_3 -alkoxy)- C_2 - C_6 -alkyl; (hydroxy)- C_2 - C_6 -alkyl; ($R^{13}R^{14}N$)- C_2 - C_6 -alkyl; ($R^{13}R^{14}NOC$)- C_1 - C_6 -alkyl; $CONR^{13}R^{14}$; $CSNR^{13}R^{14}$, or $C(NH)NR^{13}R^{14}$; and

R^{13} and R^{14} each independently of one another represent hydrogen; C_1 - C_4 -alkyl; (C_1 - C_3 -alkoxy)- C_2 - C_4 -alkyl; or (hydroxy)- C_2 - C_4 -alkyl,

and their respective salts, hydrates, alkoxides and tautomers.

12. (Amended) The combination preparation as claimed in claim 1, characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and is selected from the group consisting of 2-phenyl-substituted imidazotriazinones of the general formula




in which

R^1 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms;

R^2 represents straight-chain alkyl having up to 4 carbon atoms;

R^3 and R^4 are identical or different and represent hydrogen or represent straight-chain or branched alkenyl or alkoxy having in each case up to 8 carbon atoms, or represent a straight-chain or branched alkyl chain having up to 10 carbon atoms which is optionally interrupted by an oxygen atom and which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of trifluoromethyl, trifluoromethoxy, hydroxyl, halogen, carboxyl, benzyloxycarbonyl, straight-chain or branched alkoxycarbonyl having up to 6 carbon atoms or by radicals of the formulae $-SO_3H$, $-(A)_a-NR^7R^8$, $-O-CO-NR^7R^8$, $-S(O)_b-R^9$, $-P(O)(OR^{10})(OR^{11})$,



in which

a and b are identical or different and represent a number 0 or 1,

A represents a radical CO or SO₂,

R⁷, R^{7'}, R⁸ and R^{8'} are identical or different and represent hydrogen, or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms, a 5- to 6-membered unsaturated, partially unsaturated or saturated optionally benzo-fused heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, where the abovementioned ring systems are optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, nitro, trifluoromethyl, trifluoromethoxy, carboxyl, halogen, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms or by a group of the formula -(SO₂)_c-NR¹²R¹³,

in which

c represents a number 0 or 1,

R¹² and R¹³ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 5 carbon atoms,

or

R⁷, R^{7'}, R⁸ and R^{8'} represent straight-chain or branched alkoxy having up to 6 carbon atoms, or represent straight-chain or branched alkyl having up to 8 carbon atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, aryl having 6 to 10 carbon

atoms, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms, or by a group of the formula $-(CO)_d-NR^{14}R^{15}$,

in which

R^{14} and R^{15} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

and

d represents a number 0 or 1,

or

R^7 and R^8 or $R^{7'}$ and $R^{8'}$ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{16}$,

in which

R^{16} represents hydrogen, aryl having 6 to 10 carbon atoms, benzyl, a 5- to 7-membered aromatic or saturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, which heterocycle is optionally substituted by methyl, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl,

R^9 represents aryl having 6 to 10 carbon atoms, or represents straight-chain or branched alkyl having up to 4 carbon atoms,

R^{10} and R^{11} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

or the alkyl chain listed above under R^3/R^4 is optionally substituted by cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or by a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula $-NR^{17}$,

in which

R^{17} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl or alkoxy having in each case up to 4 carbon atoms, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl and straight-chain or branched alkoxy having up to 6 carbon atoms,

and where aryl and the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, $-SO_3H$, straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy or by a radical of the formula $-SO_2NR^{18}R^{19}$,

in which

R^{18} and R^{19} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,

or


R^3 or R^4 represent a group of the formula $-NR^{20}R^{21}$,

in which


R^{20} and R^{21} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

or

R^3 or R^4 represent adamantyl, or represent radicals of the formulae



or



or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or represent a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula - NR^{22} ,

in which


R^{22} has the meaning of R^{16} given above and is identical to or different from this meaning, or represents carboxyl, formyl or straight-chain or branched acyl having up to 5 carbon atoms,

and where cycloalkyl, aryl or the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of halogen, triazolyl, trifluoromethyl, trifluoromethoxy, carboxyl, straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, or by groups of the formulae $-SO_3H$, $-OR^{23}$, $(SO_2)_eNR^{24}R^{25}$, $-P(O)(OR^{26})(OR^{27})$,

in which

e represents a number 0 or 1,

R^{23} represents a radical of the formula



represents cycloalkyl having 3 to 7 carbon atoms, or represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by cycloalkyl having 3 to 7 carbon atoms, benzyloxy, tetrahydropyranyl, tetrahydrofuranyl, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms, carboxyl, benzyloxycarbonyl or phenyl which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkoxy having up to 4 carbon atoms, hydroxyl and halogen,

or alkyl is optionally substituted by radicals of the formulae $-\text{CO}-\text{NR}^{28}\text{R}^{29}$ or $-\text{CO}-\text{R}^{30}$,

in which

R^{28} and R^{29} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 8 carbon atoms, or

R^{28} and R^{29} together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O,

and

R^{30} represents phenyl or adamantyl,

R^{24} and R^{25} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

R^{26} and R^{27} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning

or cycloalkyl, aryl or the heterocycle are optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, carboxyl, by a 5- to 7-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O or by groups of the formula $-\text{SO}_2-\text{R}^{31}$, $\text{P}(\text{O})(\text{OR}^{32})(\text{OR}^{33})$ or $-\text{NR}^{34}\text{R}^{35}$,

in which

R^{31} is hydrogen or has the meaning of R^9 given above and is identical to or different from this meaning,

R^{32} and R^{33} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{34} and R^{35} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms, or

R^{34} and R^{35} together with the nitrogen atom form a 5- to 6-membered saturated heterocycle which may contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{36}$,

in which

R^{36} represents hydrogen, hydroxyl, straight-chain or branched alkoxycarbonyl having up to 7 carbon atoms or straight-chain or branched alkyl having up to 5 carbon atoms which is optionally substituted by hydroxyl,

or

R^3 and R^4 together with the nitrogen atom form a 5- to 7-membered unsaturated or saturated or partially unsaturated optionally benzo-fused heterocycle which may optionally contain up to 3 heteroatoms from the group consisting of S, N, O or a radical of the formula $-NR^{37}$,

in which

R^{37} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl, alkoxy or alkoxy carbonyl having in each case up to 4 carbon atoms,
or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, trifluoromethyl, carboxyl, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms or by groups of the formula $-(D)_f-NR^{38}R^{39}$, $-\text{CO}-(\text{CH}_2)_g-\text{O}-\text{CO}-R^{40}$, $-\text{CO}-(\text{CH}_2)_h-\text{OR}^{41}$ or $-\text{P}(\text{O})(\text{OR}^{42})(\text{OR}^{43})$,

in which

g and h are identical or different and represent a number 1, 2, 3 or 4,

and

f represents a number 0 or 1,

D represents a group of the formula $-\text{CO}$ or $-\text{SO}_2$,

R^{38} and R^{39} are identical or different and have the meaning of R^7 and R^8 given above,

R^{40} represents straight-chain or branched alkyl having up to 6 carbon atoms,

R^{41} represents straight-chain or branched alkyl having up to 6 carbon atoms,

R^{42} and R^{43} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

or

R^{37} represents a radical of the formula - (CO)_i-E,

in which

i represents a number 0 or 1,


E represents cycloalkyl having 3 to 7 carbon atoms or benzyl, represents aryl having 6 to 10 carbon atoms or a 5- to 6-membered aromatic heterocycle having up to 4 heteroatoms from the group consisting of S, N and O, where the ring systems listed above are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, -SO₃H, straight-chain or branched alkoxy having up to 6 carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy or by a radical of the formula -SO₂-NR⁴⁴R⁴⁵,

in which


R^{44} and R^{45} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

or

E represents radicals of the formulae



and the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally mono- to polysubstituted by identical or different substituents, if appropriate also geminally, by hydroxyl, formyl, carboxyl, straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro and groups of the formulae $-P(O)(OR^{46})(OR^{47})$,



in which

R^{46} and R^{47} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{48} is hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms,

j is a number 0 or 1,

and

R^{49} and R^{50} are identical or different and have the meaning of R^{14} and R^{15} given above,

or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, carboxyl, cycloalkyl or cycloalkyloxy having in each case 3 to 8 carbon atoms, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms or by a radical of the formula $-SO_3H$, $-NR^{51}R^{52}$ or $P(O)OR^{53}OR^{54}$,

in which

R^{51} and R^{52} are identical or different and represent hydrogen, phenyl, carboxyl, benzyl or straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

R^{53} and R^{54} are identical or different and have the meaning of R^{10} and R^{11} given above,

or the alkyl is optionally substituted by aryl having 6 to 10 carbon atoms which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, straight-chain or branched alkoxy having up to 6 carbon atoms, or by a group of the formula $-NR^{51'}R^{52'}$,

in which


$R^{51'}$ and $R^{52'}$ have the meaning of R^{51} and R^{52} given above and are identical to or different from this meaning,

or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by aryl having 6 to 10 carbon atoms or


by a 5- to 7-membered saturated, partially unsaturated or unsaturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, if appropriate also attached via an N-function, where the ring systems for their part may be substituted by hydroxyl or by straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

or

R³ and R⁴ together with the nitrogen atom form radicals of the formulae



or



R⁵ and R⁶ are identical or different and represent hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or represent straight-chain or branched alkoxy having up to 6 carbon atoms.

and their respective salts, hydrates, alkoxides and tautomers.

13. (Amended) The combination preparation as claimed in claim 1, characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and is selected from the group consisting of (a) 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one and its salts, hydrates, alkoxides and tautomers; and (b) 2-[2-ethoxy-

5-(4-ethyl-piperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f]-[1,2,4]-triazin-4-one and its salts, hydrates, alkoxides and tautomers.

14. (Amended) The combination preparation as claimed in claim 13, in that the PDE inhibitor (active compound component A) is 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one citrate or 2-[2-ethoxy-5-(4-ethylpiperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one hydrochloride trihydrate.
15. (Amended) A method for enhancing the activity of PDE inhibitors by administering an effective amount of an antilipemic.
16. (Amended) A method for the treatment of sexual dysfunction in men and women comprising administering to a host in need thereof an effective amount of the combination preparation of claim 1.
17. (Amended) The method of claim 16, characterized in that the antilipemic and the PDE inhibitor are administered either simultaneously or else successively.
18. (Amended) The method of claim 16, characterized in that the antilipemic and the PDE inhibitor are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
19. (Amended) The method of claim 16, characterized in that the antilipemic and the PDE inhibitor are present (spatially) separated, in particular as a kit-of-parts.
20. (Amended) The method of claim 16, characterized in that the antilipemic is selected from the compounds defined in claims 7 to 10.


21. (Amended) The method of claim 16, characterized in that the PDE inhibitor is selected from the compounds defined in claims 11 to 14.
22. (New) The combination preparation of claim 1 characterized in that the PDE inhibitor is a cGMP PDE inhibitor.
23. (New) The combination preparation of claim 8, characterized in that the HMG-CoA-reductase inhibitor is a statin.
24. (New) The combination preparation of claim 23, characterized in that the statin is selected from the group consisting of atorvastatin, cerivastatin, fluvastatin, lovastatin, pravastatin, itavastatin, simvastatin and (+)-(3R,5S)-bis-(7-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methanesulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid, and their respective salts, hydrates, alkoxides, esters and tautomers.
25. (New) The method of claim 15, characterized in that the PDE inhibitor is a cGMP PDE inhibitor.
26. (New) The method of claim 16, characterized in that the sexual dysfunction is erectile dysfunction.

Amended Claims (Attorney Docket No. Le A 33 965)
Version with Markings to Show Changes to Claims

1. A combination preparation, comprising as pharmaceutically active ingredients at least one active compound component A and at least one active compound component B, characterized in that the active compound component A is a PDE inhibitor, [preferably a cGMP PDE inhibitor,] and the active compound component B is an antilipemic.
2. Cancelled.
3. Cancelled.
4. Cancelled.
5. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 4], characterized in that the active compound components A and B are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
6. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 4], characterized in that the active compound components A and B are (spatially) separated, in particular as a kit-of-parts.
7. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 6], characterized in that the antilipemic (active compound component B) is selected from the group consisting of (a) HMG-CoA-reductase inhibitors; (b) squalene synthase inhibitors; (c) bile acid sequestrants; (d) fibric acid and its derivatives; (e) nicotinic acid and its analogs; (f) ω 3-fatty acids.
8. (Amended) The combination preparation as claimed in claim 7, characterized in that the antilipemic (active compound component B) is an HMG-CoA-reductase inhibitor

[and is in particular selected from the group of the statins, preferably from the group consisting of atorvastatin, cerivastatin, fluvastatin, lovastatin, pravastatin, itavastatin, simvastatin and (+)-(3R,5S)-bis-(7-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methanesulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid, and their respective salts, hydrates, alkoxides, esters and tautomers].

11. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 10], characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and is [in particular] selected from the group consisting of pyrazolopyrimidones of the general formula below



in which

- R¹ represents hydrogen; C₁-C₃-alkyl; C₁-C₃-perfluoroalkyl; or C₃-C₅-cycloalkyl;
R² denotes hydrogen; C₁-C₆-alkyl, optionally substituted by C₃-C₆-cycloalkyl; C₁-C₃-perfluoroalkyl; or C₃-C₆-cycloalkyl;
R³ is C₁-C₆-alkyl, optionally substituted by C₃-C₆-cycloalkyl; C₁-C₆-perfluoroalkyl, C₃-C₅-cycloalkyl; C₃-C₆-alkenyl; or C₃-C₆-alkinyl;
R⁴ represents C₁-C₄-alkyl, optionally substituted by OH, NR⁵R⁶, CN, CONR⁵R⁶ or CO₂R⁷; C₂-C₄-alkenyl, optionally substituted by CN, CONR⁵R⁶ or CO₂R⁷; C₂-C₄-alkanoyl, optionally substituted by NR⁵R⁶; (hydroxy)-C₂-C₄-alkyl, optionally substituted by NR⁵R⁶, (C₂-C₃-alkoxy)-C₁-C₂-alkyl, optionally substituted by OH or NR⁵R⁶, CO₂R⁷; halogen; NR⁵R⁶, NHSO₂NR⁵R⁶; NHSO₂R⁸; SO₂NR⁹R¹⁰; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, each of which is optionally substituted by methyl;

R^5 and R^6 each independently of one another denote hydrogen or C_1 - C_4 -alkyl; or together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4- $N(R^{11})$ -piperazinyl or imidazolyl group, where this group is optionally substituted by methyl or OH;

R^7 is hydrogen or C_1 - C_4 -alkyl;

R^8 represents C_1 - C_3 -alkyl, optionally substituted by NR^5R^6 ;

R^9 and R^{10} together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4- $N(R^{12})$ -piperazinyl group, where this group is optionally substituted by C_1 - C_4 -alkyl, C_1 - C_3 -alkoxy, $NR^{13}R^{14}$ or $CONR^{13}R^{14}$;

R^{11} denotes hydrogen, C_1 - C_3 -alkyl, optionally substituted by phenyl; (hydroxy)- C_2 - C_3 -alkyl; or C_1 - C_4 -alkanoyl;


R^{12} is hydrogen, C_1 - C_6 -alkyl, (C_1 - C_3 -alkoxy)- C_2 - C_6 -alkyl; (hydroxy)- C_2 - C_6 -alkyl; ($R^{13}R^{14}N$)- C_2 - C_6 -alkyl; ($R^{13}R^{14}NOC$)- C_1 - C_6 -alkyl; $CONR^{13}R^{14}$; $CSNR^{13}R^{14}$, or $C(NH)NR^{13}R^{14}$; and

R^{13} and R^{14} each independently of one another represent hydrogen; C_1 - C_4 -alkyl; (C_1 - C_3 -alkoxy)- C_2 - C_4 -alkyl; or (hydroxy)- C_2 - C_4 -alkyl,

and their respective salts, hydrates, alkoxides and tautomers.

12. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 10], characterized in that the PDE inhibitor (active compound component A) is a cGMP

PDE inhibitor and is [in particular] selected from the group consisting of 2-phenyl-substituted imidazotriazinones of the general formula





in which



R^1 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms;

R^2 represents straight-chain alkyl having up to 4 carbon atoms;

R^3 and R^4 are identical or different and represent hydrogen or represent straight-chain or branched alkenyl or alkoxy having in each case up to 8 carbon atoms, or represent a straight-chain or branched alkyl chain having up to 10 carbon atoms which is optionally interrupted by an oxygen atom and which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of trifluoromethyl, trifluoromethoxy, hydroxyl, halogen, carboxyl, benzyloxycarbonyl, straight-chain or branched alkoxycarbonyl having up to 6 carbon atoms [and/]or by radicals of the formulae $-SO_3H$, $-(A)_a-NR^7R^8$, $-O-CO-NR^7R^8$, $-S(O)_b-R^9$, $-P(O)(OR^{10})(OR^{11})$,



[and/]or



in which

a and b are identical or different and represent a number 0 or 1,

A represents a radical CO or SO₂,

R⁷, R^{7'}, R⁸ and R^{8'} are identical or different and represent hydrogen, or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms, a 5- to 6-membered unsaturated, partially unsaturated or saturated optionally benzo-fused heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, where the abovementioned ring systems are optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, nitro, trifluoromethyl, trifluoromethoxy, carboxyl, halogen, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms or by a group of the formula -(SO₂)_c-NR¹²R¹³,

in which

c represents a number 0 or 1,

R^{12} and R^{13} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 5 carbon atoms,

or

R^7 , $R^{7'}$, R^8 and $R^{8'}$ represent straight-chain or branched alkoxy having up to 6 carbon atoms, or represent straight-chain or branched alkyl having up to 8 carbon atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, aryl having 6 to 10 carbon atoms, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms, or by a group of the formula $-(CO)_d-NR^{14}R^{15}$,

in which

R^{14} and R^{15} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

and

d represents a number 0 or 1,

or

R^7 and R^8 [and/]or $R^{7'}$ and $R^{8'}$ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{16}$,

in which

R¹⁶ represents hydrogen, aryl having 6 to 10 carbon atoms, benzyl, a 5- to 7-membered aromatic or saturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, which heterocycle is optionally substituted by methyl, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl,

R⁹ represents aryl having 6 to 10 carbon atoms, or represents straight-chain or branched alkyl having up to 4 carbon atoms,

R¹⁰ and R¹¹ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

[and/]or the alkyl chain listed above under R³/R⁴ is optionally substituted by cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or by a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula -NR¹⁷,

in which

R¹⁷ represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl or alkoxy having in each case up to 4 carbon atoms, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl and straight-chain or branched alkoxy having up to 6 carbon atoms,

and where aryl and the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, $-\text{SO}_3\text{H}$, straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy [and/]or by a radical of the formula - $\text{SO}_2\text{NR}^{18}\text{R}^{19}$,

in which

R^{18} and R^{19} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,

[and/]or


R^3 or R^4 represent a group of the formula $-\text{NR}^{20}\text{R}^{21}$,

in which


R^{20} and R^{21} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

[and/]or

R^3 or R^4 represent adamantyl, or represent radicals of the formulae



or



or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or represent a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula -NR²²,

in which


R²² has the meaning of R¹⁶ given above and is identical to or different from this meaning, or represents carboxyl, formyl or straight-chain or branched acyl having up to 5 carbon atoms,

and where cycloalkyl, aryl [and/]or the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of halogen, triazolyl, trifluoromethyl, trifluoromethoxy, carboxyl, straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, [and/]or by groups of the formulae -SO₃H, -OR²³, (SO₂)_eNR²⁴R²⁵, -P(O)(OR²⁶)(OR²⁷),

in which

e represents a number 0 or 1,

R²³ represents a radical of the formula



represents cycloalkyl having 3 to 7 carbon atoms, or
represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by cycloalkyl having 3 to 7 carbon atoms, benzyloxy, tetrahydropyranyl, tetrahydrofuranyl, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms, carboxyl, benzyloxycarbonyl or phenyl which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkoxy having up to 4 carbon atoms, hydroxyl and halogen,

[and/]or alkyl is optionally substituted by radicals of the formulae -CO-NR²⁸R²⁹ or -CO-R³⁰,

in which

R²⁸ and R²⁹ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 8 carbon atoms, or

R²⁸ and R²⁹ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O,

and

R³⁰ represents phenyl or adamantyl,

R^{24} and R^{25} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

R^{26} and R^{27} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning

[and/]or cycloalkyl, aryl [and/]or the heterocycle are optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, carboxyl, by a 5- to 7-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O or by groups of the formula - SO_2-R^{31} , $P(O)(OR^{32})(OR^{33})$ or $-NR^{34}R^{35}$,

in which

R^{31} is hydrogen or has the meaning of R^9 given above and is identical to or different from this meaning,

R^{32} and R^{33} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{34} and R^{35} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms, or

R^{34} and R^{35} together with the nitrogen atom form a 5- to 6-membered saturated heterocycle which may contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{36}$,

in which

R^{36} represents hydrogen, hydroxyl, straight-chain or branched alkoxy carbonyl having up to 7 carbon atoms or straight-chain or branched alkyl having up to 5 carbon atoms which is optionally substituted by hydroxyl,

or

R^3 and R^4 together with the nitrogen atom form a 5- to 7-membered unsaturated or saturated or partially unsaturated optionally benzo-fused heterocycle which may optionally contain up to 3 heteroatoms from the group consisting of S, N, O or a radical of the formula $-NR^{37}$,

in which

R^{37} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl, alkoxy or alkoxy carbonyl having in each case up to 4 carbon atoms,
or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, trifluoromethyl, carboxyl, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms or by groups of the formula $-(D)_f-NR^{38}R^{39}$, $-\text{CO}-(\text{CH}_2)_g-\text{O}-\text{CO}-R^{40}$, $-\text{CO}-(\text{CH}_2)_h-\text{OR}^{41}$ or $-\text{P}(\text{O})(\text{OR}^{42})(\text{OR}^{43})$,

in which

g and h are identical or different and represent a number 1, 2, 3 or 4,

and

f represents a number 0 or 1,

D represents a group of the formula -CO or -SO₂,

R³⁸ and R³⁹ are identical or different and have the meaning of R⁷ and R⁸ given above,

R⁴⁰ represents straight-chain or branched alkyl having up to 6 carbon atoms,

R⁴¹ represents straight-chain or branched alkyl having up to 6 carbon atoms,

R⁴² and R⁴³ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

or

R³⁷ represents a radical of the formula - (CO)_i-E,

in which

i represents a number 0 or 1,

E represents cycloalkyl having 3 to 7 carbon atoms or benzyl,
represents aryl having 6 to 10 carbon atoms or a 5- to 6-membered aromatic heterocycle having up to 4 heteroatoms from the group consisting of S, N and O, where the ring systems listed above are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, -SO₃H,


straight-chain or branched alkoxy having up to 6 carbon atoms,
hydroxyl, trifluoromethyl, trifluoromethoxy or by a radical of the
formula $-\text{SO}_2-\text{NR}^{44}\text{R}^{45}$,

in which


R^{44} and R^{45} have the meaning of R^{18} and R^{19} given above and are
identical to or different from this meaning,

or

E represents radicals of the formulae



and the heterocycle listed under R^3 and R^4 , which is formed together with the
nitrogen atom, is optionally mono- to polysubstituted by identical or different
substituents, if appropriate also geminally, by hydroxyl, formyl, carboxyl,
straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6
carbon atoms, nitro and groups of the formulae $-\text{P}(\text{O})(\text{OR}^{46})(\text{OR}^{47})$,



in which

R^{46} and R^{47} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{48} is hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms,

j is a number 0 or 1,

and

R^{49} and R^{50} are identical or different and have the meaning of R^{14} and R^{15} given above,

[and/]or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, carboxyl, cycloalkyl or cycloalkyloxy having in each case 3 to 8 carbon atoms, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms or by a radical of the formula $-\text{SO}_3\text{H}$, $-\text{NR}^{51}\text{R}^{52}$ or $\text{P}(\text{O})\text{OR}^{53}\text{OR}^{54}$,

in which

R^{51} and R^{52} are identical or different and represent hydrogen, phenyl, carboxyl, benzyl or straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

R^{53} and R^{54} are identical or different and have the meaning of R^{10} and R^{11} given above,

[and/]or the alkyl is optionally substituted by aryl having 6 to 10 carbon atoms which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, straight-chain or branched alkoxy having up to 6 carbon atoms, or by a group of the formula -NR^{51'}R^{52'},




in which


R^{51'} and R^{52'} have the meaning of R⁵¹ and R⁵² given above and are identical to or different from this meaning,

[and/]or the heterocycle listed under R³ and R⁴, which is formed together with the nitrogen atom, is optionally substituted by aryl having 6 to 10 carbon atoms or by a 5- to 7-membered saturated, partially unsaturated or unsaturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, if appropriate also attached via an N-function, where the ring systems for their part may be substituted by hydroxyl or by straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

or

R³ and R⁴ together with the nitrogen atom form radicals of the formulae





R⁵ and R⁶ are identical or different and represent hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or represent straight-chain or branched alkoxy having up to 6 carbon atoms.

and their respective salts, hydrates, alkoxides and tautomers.

13. (Amended) The combination preparation as claimed in [any of] claim[s] 1 [to 10], characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and [in particular] is selected from the group consisting of (a) 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one [(sildenafil)] and its salts, hydrates, alkoxides and tautomers; and (b) 2-[2-ethoxy-5-(4-ethyl-piperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f]-[1,2,4]-triazin-4-one and its salts, hydrates, alkoxides and tautomers.
14. (Amended) The combination preparation as claimed in claim 13, in that the PDE inhibitor (active compound component A) is 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one citrate [(sildenafil citrate, ViagraTM)] or 2-[2-ethoxy-5-(4-ethylpiperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one hydrochloride trihydrate.
15. (Amended) [The use of antilipemics] A method for enhancing the activity of PDE inhibitors [, in particular cGMP PDE inhibitors] by administering an effective amount of an antilipemic.

16. (Amended) [The use as claimed in claim 15 in] A method for the treatment of sexual dysfunction in men and women comprising administering to a host in need thereof an effective amount of the combination preparation of claim 1 [, in particular in the treatment of erectile dysfunction].
17. (Amended) The [use as claimed in claim 15 or] method of claim 16, characterized in that the antilipemic and the PDE inhibitor are [used] administered either simultaneously or else successively.
18. (Amended) The [use as claimed in any of claims 15 to 17] method of claim 16, characterized in that the antilipemic and the PDE inhibitor are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
19. (Amended) The [use as claimed in any of claims 15 to 17] method of claim 16, characterized in that the antilipemic and the PDE inhibitor are present (spatially) separated, in particular as a kit-of-parts.
20. (Amended) The [use as claimed in any of claims 15 to 19] method of claim 16, characterized in that the antilipemic is selected from the compounds defined in claims 7 to 10.

Novel combination for treating sexual dysfunction

The present invention relates to the area of sexual dysfunction in men and women.

- 5 The present invention relates in particular to a novel combination preparation for treating sexual dysfunction in men and women, in particular erectile dysfunction. The present invention thus also provides a novel combination therapy for the treatment of sexual dysfunction in men and women.
- 10 The present invention furthermore relates to the use of antilipemics for enhancing the activity of phosphodiesterase inhibitors (hereinbelow synonymously also referred to as "phosphodiesterase inhibitors" or "PDE inhibitors") for treatment of sexual dysfunction.
- 15 From the prior art it is known that phosphodiesterase inhibitors – in particular those of subtype V, which are synonymously also referred to as "cGMP PDE inhibitors" – are suitable for treating sexual dysfunction, in particular for treating erectile dysfunction (see, for example, *Molecular Pharmacology*, 1999, **56**, pages 124-130; *Am. J. Physiol.*, Vol. 264, February 1993, pages H419-H422; *The Journal of*
20 *Urology*, Vol. 147, pages 1650-1655 (June 1992); *The New England Journal of Medicine*, Vol. 326(2), pages 90-94 (9 January 1992); *International Journal of Impotence Research*, **4**, Suppl. 2, page 11 (1992); *Drugs, News and Perspectives*, **6**(3), pages 150-156 (April 1993); *Physiological Reviews* **75**, pages 191-236 (1995); *Int. J. of Impotence* **9**, pages 17-26 (1997) and *TIPS Reviews*, Vol. 11, pages 150-155
25 (April 1990)).

Concerning the nomenclature of PDE inhibitors, reference is made to *Beavo* and *Reifsnnyder* in *Trends in Pharmacol. Sci.* 1990. **11**, pages 150-155 and to the article *TIPS Reviews*, Vol. 11, pages 150-155 (April 1990).

30

The reason for the effectiveness of PDE inhibitors, in particular cGMP PDE inhibitors (PDE V inhibitors), in the treatment of sexual, preferably erectile, dysfunction is that the neurotransmitter nitrogen monoxide NO, which is generated

by the body following sexual stimulation, activates guanylate cyclase, which in turn converts GTP into cGMP, which for its part may then cause a relaxation of the corpus cavernosum, without the cGMP, which is responsible for the relaxation of the corpus cavernosum, being hydrolyzed by the enzyme phosphodiesterase V (PDE V) to give 5'GMP, since the activity of the enzyme PDE V is inhibited by the corresponding inhibitor. This mechanism is illustrated in the attached **Fig. 1**.

However, it has now been found that, in some patients, therapy of erectile dysfunction using PDE inhibitors, in particular cGMP PDE inhibitors, is unsuccessful or of only limited success. This group of patients comprises in particular patients having a disturbed endothelium function and/or metabolic disorders, such as, for example, hyperlipidemia (for example hypercholesterolemia), arteriosclerosis, diabetes (in particular of the diabetes mellitus type), and also heavy smokers and elderly patients. In these patients, the therapy of erectile dysfunction using customary doses results only in a considerably reduced effect, compared to other groups of patients which undergo therapy for erectile dysfunction but do not suffer from the abovementioned metabolic disorders.

However, it is exactly the abovementioned patient group having disturbed endothelium function and/or the abovementioned metabolic disorders which suffers more frequently than average from sexual dysfunction, in particular erectile dysfunction, which renders the customary treatment of this dysfunction with PDE inhibitors more difficult. To achieve a therapeutic effect, considerably higher doses of PDE inhibitors have to be administered to these problem patients, compared to other patient groups suffering from erectile dysfunction, but not from the metabolic disorders mentioned above. This, however, has – in addition to higher costs – the essential disadvantage that the side effects associated with the PDE inhibitor therapy are also increased by the factor of the higher dosage. These side effects include, for example, disturbed vision, in particular disturbed color vision and color perception, headaches and muscle aches. Side effects on the cardiovascular system, for example a lowering of the blood pressure, are also possible.

Surprisingly, the applicant has now found that the activity of PDE inhibitors, in

particular PDE V inhibitors, in the therapy of sexual, preferably erectile, dysfunction – in particular in the groups of problem patients mentioned above – can be increased by combined administration of the PDE inhibitors with an antilipemic. In this manner, it is possible to avoid the disadvantages described above which are encountered in the customary treatment of erectile dysfunction when using PDE inhibitors alone, in particular in the problem patients mentioned above.

Accordingly, the present invention relates to a combination preparation comprising

- as active compound component A at least one PDE inhibitor, preferably a PDE V inhibitor (cGMP PDE inhibitor); and
- as active compound component B at least one antilipemic.

The combination preparation according to the invention is particularly suitable for therapy of sexual dysfunction, i.e. therapy of erectile dysfunction in men or sexual dysfunction in women.

At the same time, the combination preparation according to the invention allows a cotherapy of disturbed endothelium function (for example in elderly patients or heavy smokers) and/or a metabolic disorder, such as, for example, hyperlipidemia (for example hypercholesterolemia), arteriosclerosis or diabetes (in particular of the diabetes mellitus type).

The term “combination preparation”, as used for the purpose of the present invention, means that the two active compound components A and B can be used either simultaneously or else successively (i.e. separately).

Thus, according to the invention, the term “combination preparation” includes the ingredients A and B either in a functional unit, i.e. as a true combination (for example as a mixture, mix or blend), or else (spatially) separated, i.e. as a “kit-of-parts”.

Accordingly, the present invention also provides the use of antilipemics for enhancing the activity of PDE inhibitors (in particular PDE V inhibitors) in the

therapy of sexual dysfunction, in particular erectile dysfunction.

The present invention furthermore provides a combination therapy for sexual dysfunction, in particular for erectile dysfunction, using a combination preparation
5 which comprises at least one PDE inhibitor (in particular a PDE V inhibitor) and at least one antilipemic.

At the same time, the combination therapy according to the invention also allows a cotherapy of a disturbed endothelium function (for example in elderly patients or
10 heavy smokers) and/or a metabolic disorder, such as, for example, hyperlipidemia (for example hypercholesterolemia), arteriosclerosis or diabetes (in particular of the diabetes mellitus type).

Using the combination according to the invention of PDE inhibitors and antilipemics
15 – i.e., in other words, by applying the combination therapy according to the invention of sexual dysfunction using a combination of PDE inhibitor and antilipemic – it is possible to reduce the therapeutically required doses of PDE inhibitors in the abovementioned problem patients to customary doses like those administered to other patients suffering from erectile dysfunction but not from disturbed endothelium
20 function or a metabolic disorder. However, according to the invention, it is also possible to reduce, by combined administration with an antilipemic, the dose of PDE inhibitor administered for the therapy of erectile dysfunction in patients who do not suffer from a specific metabolic disorder but who do suffer (for example because of old age) from erectile dysfunction.

25

As mentioned above, the combination according to the invention can be administered, i.e. the combination therapy according to the invention can be applied, by simultaneous administration of active compound components A and B. Here, the active compound components A and B can, as described above, be present either in a
30 functional unit (i.e. as a true combination, such as, for example, as a mixture, a mix or a blend) or else (spatially) separated (i.e. as a “kit” or a “kit-of-parts”).

According to a preferred embodiment of the present invention, the active compound

components A and B are administered separately, in particular successively.

This can take place, for example, by administering a daily dose of the antilipemic even a few days (for example about 1 week or else only 1-4 days) before
5 administering the PDE inhibitor.

It is also possible to administer the PDE inhibitor in an ongoing therapy with antilipemics. Thus, according to the present invention it is possible, for example, to observe better successes in the therapy of erectile dysfunction using PDE inhibitors
10 in men suffering from severe hypercholesterolemia in which the elevated cholesterol levels are already treated on a permanent basis using antilipemics.

However, surprisingly, similarly good therapeutic results are also observed in cases where the antilipemic is administered only a short time (for example a few days)
15 before the administration of the PDE inhibitor. This is surprising and indicates an unforeseeable synergistic effect of the combination according to the invention, since in the patients treated in this manner, suffering, for example, from arteriosclerosis, there are not yet any changes in their symptoms, owing to the short duration of the therapy with the antilipemic.

20 According to a preferred embodiment of the present invention, the active compound components A and B of the combination preparation according to the invention are thus administered successively, with the antilipemic preferably being administered first, i.e. before the administration of the PDE inhibitor. This can be achieved by
25 simply administering the antilipemic a short time before the administration of the PDE inhibitor, i.e. several times over a number of days or else only once a few hours beforehand, or else by administering the PDE inhibitor in an ongoing therapy with an antilipemic. Thus, in the latter case, the administration of the antilipemic can be continued both before and concomitantly with the administration of the PDE
30 inhibitor.

Without wishing to adhere to a certain theory, the improved PDE-inhibitory action of the PDE inhibitor by simultaneous or successive or concomitant administration of

antilipemics can probably be explained by the fact that antilipemics improve the disturbed endothelium function by generating nitrogen monoxide (NO) (*Current Opinion in Lipidology*, 1997, Vol. 8, pages 362-368 and *Circulation* 1998, **97**, pages 1129-1135). As a neurotransmitter, nitrogen monoxide in turn is an important physiological factor for an erection to take place, since it increases the cGMP concentration, which finally results in a relaxation of the corpus cavernosum (*International Journal of Impotence Research* 1999, **11**, pages 123-132 and 159-165).

10 According to the present invention, the antilipemic can be selected from the group consisting of:

- HMG-CoA-reductase inhibitors
- squalene synthase inhibitors,
- bile acid absorption inhibitors (also referred to as "bile acid anion exchangers" or
- 15 bile acid sequestrants),
- fibric acid and its derivatives,
- nicotinic acids and its analogs and also
- ω 3-fatty acids.

20 For further details about the antilipemics mentioned above, reference is in this context made to the article by Gilbert R. Thompson & Rissitaza P. Naoumova "New prospects for lipid-lowering drugs" in *Exp. Opin. Invest. Drugs* (1998), **7**(5), pages 715-727, the entire content of which is hereby expressly incorporated by way of reference.

25

Among the antilipemics mentioned above, preference according to the invention is given to the HMG-CoA-reductase inhibitors. Here, the abbreviation "HMG-CoA" denotes "3-hydroxymethylglutaryl-coenzyme A".

30 Among the HMG-CoA-reductase inhibitors, in turn, preference according to the invention is given, in particular, to the substance class of the vastatins – which, for the sake of simplicity, are in most cases referred to in the literature simply as

“statins”.

Among the statins, in turn, particular preference according to the invention is given to

- 5 • atorvastatin (commercially available under the name Lipitor® from Parke-Davis);
- cerivastatin (commercially available under the name Lipobay® or Baycol® from Bayer);
- fluvastatin (commercially available under the name Lescol® from Novartis);
- lovastatin (commercially available under the name Mevacor® from Merck);
- 10 • pravastatin (commercially available under the name Lipostat® from Bristol-Myers Squibb);
- simvastatin (commercially available under the name Zocor® from Merck);
- itavastatin (also called “nisvastatin”; NK-104; systematic name:
[S-[R*,S*-(E)]]-7-[2-cyclopropyl-4-(4-fluorophenyl)-3-quinolinyl]-
15 3,5-dihydroxy-6-heptenoic acid);
- dalvastatin;
- mevastatin;
- dihydrocompactin;
- compactin; and
- 20 • (+)-(3R,5S)-bis-(7-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methanesulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid;
and their respective salts, hydrates, alkoxides, esters and tautomers,
and among these with very particular preference atorvastatin, cerivastatin,
fluvastatin, lovastatin, pravastatin, itavastatin, simvastatin and (+)-(3R,5S)-bis-(7-(4-
25 (4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methanesulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid and their respective salts, hydrates, alkoxides,
esters and tautomers.

Among these, in turn, very particular preference is given to cerivastatin and
30 atorvastatin and their respective salts, hydrates, alkoxides, esters and tautomers.

For further details about the statins mentioned above, reference is made to the

discourse in *Drugs of the Future* 1994, 19(6), pages 537-541 and 1995, 20(6), page 611 and 1996, 21(6), page 642, the respective content of which is included herein in its entirety by way of reference.

5 For the purpose of the present invention, the term "salt" refers in each case to physiologically acceptable salts of the compounds in question: these can, for example, be salts with mineral acids, carboxylic acids or sulfonic acids, in particular with hydrochloric acid, hydrobromic acid, sulfuric acid, phosphoric acid, methanesulfonic acid, ethanesulfonic acid, toluenesulfonic acid, benzenesulfonic
10 acid, naphthalenedisulfonic acid, acetic acid, propionic acid, lactic acid, tartaric acid, citric acid, fumaric acid, maleic acid or benzoic acid, or else mixed salts thereof. However, the salts can also be salts with customary bases, such as, for example, alkali metal salts (for example sodium or potassium salts), alkaline earth metal salts (for example calcium or magnesium salts) or ammonium salts, derived from
15 ammonia or organic amines, such as, for example, diethylamine, triethylamine, ethyldiisopropylamine, procaine, dibenzylamine, N-methylmorpholine, dihydro-abietylamine, 1-phenamine or methyl-piperidine, and also mixed salts thereof.

Examples of statin salts which can be used according to the invention are
20 fluindostatin (the monosodium salt of fluvastatin); the monopotassium salt and the calcium salt of itavastatin; and the calcium salt of (+)-(3R,5S)-bis-(7-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methane-sulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid ("ZD 4522" or "S 4522" from Shionogi and AstraZeneca, respectively). Further examples of statin salts which can be used
25 according to the invention are the monosodium and the monopotassium salts and also the calcium salts of cerivastatin, atorvastatin and pravastatin.

Further preferred HMG-CoA-reductase inhibitors are described in EP-A-0 325 130 and EP-A-0-491 226, both in the name of Bayer AG, the content of which is hereby
30 included by way of reference. EP-A-0 325 130 provides substituted pyridines, and EP-A-0-491 226 describes substituted pyridyldihydroxyheptenoic acid derivatives and their salts, and among these in particular cerivastatin, which is particularly preferred according to the invention (claim 6 of EP-A-0-491 226).

Preference according to the invention is also given to the statins mentioned in WO-A-99/11263, the disclosure of which is included by way of reference.

- 5 Preference according to the invention is likewise given to the HMG-CoA-reductase inhibitors mentioned in the publication *Bioorganic & Medicinal Chemistry*, Vol. 5, No. 2, pages 437-444 (1997), the disclosure of which is hereby included in its entirety by way of reference.
- 10 A further review of HMG-CoA-reductase inhibitors can be found in *Pharmazie in unserer Zeit*, Vol. 28, No. 3, pages 147-1152 (1999).

- Among the bile acid sequestrants mentioned above, preference according to the invention is given to cholestyramine (commercially available under the name
- 15 Questran[®] from Bristol-Myers Squibb) and colestipol (commercially available under the name Colestid[®] from Pharmacia & Upjohn) (see also *Exp. Opin. Invest. Drugs* (1998), 7(5), pages 715-727).


- Among the fibric acid derivatives mentioned above, preference according to the
- 20 invention is given to ciprofibrate (commercially available under the name Modalim[®] from Sanofi Winthrop), fenofibrate (commercially available under the name Lipantil[®] from Fournier), gemfibrozil (commercially available under the name Lopid[®] from Parke-Davis), bezafibrate and chlofibrate (see also *Exp. Opin. Invest. Drugs* (1998), 7(5), pages 715-727).

- 25 Among the nicotinic acid analogs mentioned above, preference according to the invention is given to acipimox (commercially available under the name Olbetam[®] from Pharmacia & Upjohn) (see also *Exp. Opin. Invest. Drugs* (1998), 7(5), pages 715-727).

- 30 Among the ω 3-fatty acids mentioned above, preference according to the invention is given to Maxepa (distributed by Seven Seas) (see also *Exp. Opin. Invest. Drugs* (1998), 7(5), pages 715-727).

Among the phosphodiesterase inhibitors, preference according to the invention is given in particular to cGMP PDE inhibitors. Among these, in turn, preference is given in particular to those which can be administered orally and, at the same time, also have good activity following oral administration.

Examples of cGMP PDE inhibitors which can be used and are preferred according to the invention are in particular the pyrazolopyrimidones described in EP-A-0 463 756, EP-A-0 526 004, WO-A-94/28902 and EP-B-0 702 555, the respective content of which is hereby included by way of reference. These are in particular compounds of the general formula below



in which

- 15 R^1 represents: hydrogen; C_1 - C_3 -alkyl; C_1 - C_3 -perfluoroalkyl; or C_3 - C_5 -cycloalkyl;
 R^2 denotes: hydrogen; C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl;
 C_1 - C_3 -perfluoroalkyl; or C_3 - C_6 -cycloalkyl;
 R^3 is: C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl; C_1 - C_6 -perfluoroalkyl;
 C_3 - C_5 -cycloalkyl; C_3 - C_6 -alkenyl; or C_3 - C_6 -alkinyl;
- 20 R^4 represents: C_1 - C_4 -alkyl, optionally substituted by OH, NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkenyl, optionally substituted by CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkanoyl, optionally substituted by NR^5R^6 ; (hydroxy)- C_2 - C_4 -alkyl, optionally substituted by NR^5R^6 , (C_2 - C_3 -alkoxy)- C_1 - C_2 -alkyl, optionally substituted by OH or NR^5R^6 , CO_2R^7 ; halogen; NR^5R^6 , $NHSO_2NR^5R^6$, $NHSO_2R^8$, $SO_2NR^9R^{10}$; or phenyl,
- 25 pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, each of which is optionally substituted by methyl;
- R^5 and R^6 each independently of one another denote hydrogen or C_1 - C_4 -alkyl; or together with the nitrogen atom to which they are attached form a pyrrolidinyl,

piperidino, morpholino, 4-N(R¹¹)-piperazinyl or imidazolyl group, where this group is optionally substituted by methyl or OH;

R⁷ is hydrogen or C₁-C₄-alkyl;

R⁸ represents C₁-C₃-alkyl, optionally substituted by NR⁵R⁶;

- 5 R⁹ and R¹⁰ together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4-N(R¹²)-piperazinyl group, where this group is optionally substituted by C₁-C₄-alkyl, C₁-C₃-alkoxy, NR¹³R¹⁴ or CONR¹³R¹⁴;

R¹¹ denotes hydrogen, C₁-C₃-alkyl, optionally substituted by phenyl; (hydroxy)-C₂-C₃-alkyl; or C₁-C₄-alkanoyl;

10


R¹² is hydrogen, C₁-C₆-alkyl, (C₁-C₃-alkoxy)-C₂-C₆-alkyl; (hydroxy)-C₂-C₆-alkyl; (R¹³R¹⁴N)-C₂-C₆-alkyl; (R¹³R¹⁴NOC)-C₁-C₆-alkyl; CONR¹³R¹⁴; CSNR¹³R¹⁴, or C(NH)NR¹³R¹⁴; and

- 15 R¹³ and R¹⁴ each independently of one another represent hydrogen; C₁-C₄-alkyl; (C₁-C₃-alkoxy)-C₂-C₄-alkyl; or (hydroxy)-C₂-C₄-alkyl,

and their respective salts, hydrates, alkoxides and tautomers.

- 20 Among these, very particular preference according to the invention is given to 5-[3-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-6,7-dihydro-1H-pyrazolo-[4,3-d]-pyrimidin-7-one (other name: 5-[2-ethoxy-5-(4-methyl-1-piperazinyl-sulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one, also known as "sildenafil") and its salts, thus, for example,
- 25 in particular the citrate salt, which is commercially available under the name ViagraTM (see in particular example 12 and compound 3 of claim 3 of EP-A-0 463 756 and also claim 6 of EP-B-0 702 555).

- 30 cGMP PDE inhibitors which are likewise preferred according to the invention are the compounds described in WO-A-99/24433, the content of which is hereby included in its entirety by way of reference. The compounds in question are 2-phenyl-substituted imidazotriazinones of the general formula





in which


R^1 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms;

R^2 represents straight-chain alkyl having up to 4 carbon atoms;

R^3 and R^4 are identical or different and represent hydrogen or represent straight-chain or branched alkenyl or alkoxy having in each case up to 8 carbon atoms, or

represent a straight-chain or branched alkyl chain having up to 10 carbon atoms which is optionally interrupted by an oxygen atom and which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of trifluoromethyl, trifluoromethoxy, hydroxyl, halogen, carboxyl, benzyloxycarbonyl, straight-chain or branched alkoxycarbonyl having up to 6 carbon atoms and/or by radicals of the formulae $-\text{SO}_3\text{H}$, $-(\text{A})_a-\text{NR}^7\text{R}^8$, $-\text{O}-\text{CO}-\text{NR}^7\text{R}^8$, $-\text{S}(\text{O})_b-\text{R}^9$, $-\text{P}(\text{O})(\text{OR}^{10})(\text{OR}^{11})$,





in which

a and b are identical or different and represent a number 0 or 1,

5

A represents a radical CO or SO₂,

R⁷, R^{7'}, R⁸ and R^{8'} are identical or different and represent hydrogen, or
 represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10
 carbon atoms, a 5- to 6-membered unsaturated, partially unsaturated
 or saturated optionally benzo-fused heterocycle having up to 3
 heteroatoms from the group consisting of S, N and O, where the
 abovementioned ring systems are optionally mono- to polysubstituted
 by identical or different substituents from the group consisting of
 hydroxyl, nitro, trifluoromethyl, trifluoromethoxy, carboxyl, halogen,
 straight-chain or branched alkoxy or alkoxycarbonyl having in each
 case up to 6 carbon atoms or by a group of the formula
 -(SO₂)_c-NR¹²R¹³,

10

15

20

in which

c represents a number 0 or 1,

R¹² and R¹³ are identical or different and represent hydrogen or
 straight-chain or branched alkyl having up to 5 carbon atoms,

25

or

R^7 , $R^{7'}$, R^8 and $R^{8'}$ represent straight-chain or branched alkoxy having up to 6 carbon atoms, or

represent straight-chain or branched alkyl having up to 8 carbon atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, aryl having 6 to 10 carbon atoms, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms, or by a group of the formula $-(CO)_d-NR^{14}R^{15}$,

in which

R^{14} and R^{15} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

and

d represents a number 0 or 1,

or

R^7 and R^8 and/or $R^{7'}$ and $R^{8'}$ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{16}$,

in which

R^{16} represents hydrogen, aryl having 6 to 10 carbon atoms, benzyl, a 5- to 7-membered aromatic or saturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, which heterocycle is optionally substituted by methyl, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl,

R^9 represents aryl having 6 to 10 carbon atoms, or
represents straight-chain or branched alkyl having up to 4 carbon
atoms,

R^{10} and R^{11} are identical or different and represent hydrogen or straight-chain
or branched alkyl having up to 4 carbon atoms,

and/or the alkyl chain listed above under R^3/R^4 is optionally substituted by
cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or
by a 5- to 7-membered partially unsaturated, saturated or unsaturated
optionally benzo-fused heterocycle which may contain up to 4 heteroatoms
from the group consisting of S, N; O or a radical of the formula $-NR^{17}$,

in which

R^{17} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain
or branched acyl or alkoxy having in each case up to 4 carbon atoms,
or represents straight-chain or branched alkyl having up to 6 carbon
atoms which is optionally mono- to polysubstituted by identical or
different substituents from the group consisting of hydroxyl and
straight-chain or branched alkoxy having up to 6 carbon atoms,

and where aryl and the heterocycle are optionally mono- to polysubstituted by
identical or different substituents from the group consisting of nitro, halogen,
 $-SO_3H$, straight-chain or branched alkyl or alkoxy having in each case up to 6
carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy and/or by a radical
of the formula $-SO_2NR^{18}R^{19}$,

in which

R^{18} and R^{19} are identical or different and represent hydrogen or straight-chain
or branched alkyl having up to 6 carbon atoms,

and/or

R^3 or R^4 represent a group of the formula $-NR^{20}R^{21}$,

5


in which

R^{20} and R^{21} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,


10

and/or

R^3 or R^4 represent adamantyl, or
represent radicals of the formulae



or



15

or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or represent a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula $-NR^{22}$,

20

in which

R^{22} has the meaning of R^{16} given above and is identical to or different from this meaning, or

25


represents carboxyl, formyl or straight-chain or branched acyl having up to 5 carbon atoms,

and where cycloalkyl, aryl and/or the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of halogen, triazolyl, trifluoromethyl, trifluoromethoxy, carboxyl, straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro, and/or by groups of the formulae $-\text{SO}_3\text{H}$, $-\text{OR}^{23}$, $(\text{SO}_2)_e\text{NR}^{24}\text{R}^{25}$, $-\text{P}(\text{O})(\text{OR}^{26})(\text{OR}^{27})$,

in which

e represents a number 0 or 1,

R^{23} represents a radical of the formula



represents cycloalkyl having 3 to 7 carbon atoms, or

represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by cycloalkyl having 3 to 7 carbon atoms, benzyloxy, tetrahydropyranyl, tetrahydrofuranyl, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms, carboxyl, benzyloxycarbonyl or phenyl which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of straight-chain or branched alkoxy having up to 4 carbon atoms, hydroxyl and halogen,

and/or alkyl is optionally substituted by radicals of the formulae $-\text{CO}-\text{NR}^{28}\text{R}^{29}$ or $-\text{CO}-\text{R}^{30}$,

in which

R^{28} and R^{29} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 8 carbon atoms, or

5 R^{28} and R^{29} together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O,

and

10

R^{30} represents phenyl or adamantyl,

R^{24} and R^{25} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

15

R^{26} and R^{27} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning

20

and/or cycloalkyl, aryl and/or the heterocycle are optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, carboxyl, by a 5- to 7-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O or by groups of the formula $-SO_2-R^{31}$, $P(O)(OR^{32})(OR^{33})$ or $-NR^{34}R^{35}$,

25

in which

R^{31} is hydrogen or has the meaning of R^9 given above and is identical to or different from this meaning,

30

R^{32} and R^{33} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{34} and R^{35} are identical or different and represent hydrogen or straight-chain

or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms, or

5 R^{34} and R^{35} together with the nitrogen atom form a 5- to 6-membered saturated heterocycle which may contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{36}$,

in which

10

R^{36} represents hydrogen, hydroxyl, straight-chain or branched alkoxy-carbonyl having up to 7 carbon atoms or straight-chain or branched alkyl having up to 5 carbon atoms which is optionally substituted by hydroxyl,

15

or

R^3 and R^4 together with the nitrogen atom form a 5- to 7-membered unsaturated or saturated or partially unsaturated optionally benzo-fused heterocycle which
20 may optionally contain up to 3 heteroatoms from the group consisting of S, N, O or a radical of the formula $-NR^{37}$,

in which

25 R^{37} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl, alkoxy or alkoxy-carbonyl having in each case up to 4 carbon atoms,

or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or
30 different substituents from the group consisting of hydroxyl, trifluoromethyl, carboxyl, straight-chain or branched alkoxy or alkoxy-carbonyl having in each case up to 6 carbon atoms or by groups of the formula $-(D)_f-NR^{38}R^{39}$, $-CO-(CH_2)_g-O-CO-R^{40}$,

- 20 -

$-\text{CO}-(\text{CH}_2)_h-\text{OR}^{41}$ or $-\text{P}(\text{O})(\text{OR}^{42})(\text{OR}^{43})$,

in which

5 g and h are identical or different and represent a number 1, 2, 3 or 4,

and

10 f represents a number 0 or 1,

D represents a group of the formula $-\text{CO}$ or $-\text{SO}_2$,

15 R^{38} and R^{39} are identical or different and have the meaning of R^7 and R^8 given above,

R^{40} represents straight-chain or branched alkyl having up to 6 carbon atoms,

20 R^{41} represents straight-chain or branched alkyl having up to 6 carbon atoms,

R^{42} and R^{43} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

25 or

R^{37} represents a radical of the formula $-(\text{CO})_i-\text{E}$,

in which

30

i represents a number 0 or 1,

E represents cycloalkyl having 3 to 7 carbon atoms or benzyl,

- 21 -



represents aryl having 6 to 10 carbon atoms or a 5- to 6-membered aromatic heterocycle having up to 4 heteroatoms from the group consisting of S, N and O, where the ring systems listed above are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, $-\text{SO}_3\text{H}$, straight-chain or branched alkoxy having up to 6 carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy or by a radical of the formula $-\text{SO}_2-\text{NR}^{44}\text{R}^{45}$,

in which


R^{44} and R^{45} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

or



E represents radicals of the formulae



or



and the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally mono- to polysubstituted by identical or different substituents, if appropriate also geminally, by hydroxyl, formyl, carboxyl, straight-chain or branched acyl or alkoxycarbonyl having in each case up to 6 carbon atoms, nitro and groups of the formulae $-\text{P}(\text{O})(\text{OR}^{46})(\text{OR}^{47})$,



in which

R^{46} and R^{47} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{48} is hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms,

j is a number 0 or 1,

and

R^{49} and R^{50} are identical or different and have the meaning of R^{14} and R^{15} given above,

and/or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, carboxyl, cycloalkyl or cycloalkyloxy having in each case 3 to 8 carbon atoms, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms or by a radical of the formula $-SO_3H$, $-NR^{51}R^{52}$ or $P(O)OR^{53}OR^{54}$,

in which

R^{51} and R^{52} are identical or different and represent hydrogen, phenyl, carboxyl, benzyl or straight-chain or branched alkyl or alkoxy having

in each case up to 6 carbon atoms,

R^{53} and R^{54} are identical or different and have the meaning of R^{10} and R^{11} given above,

and/or the alkyl is optionally substituted by aryl having 6 to 10 carbon atoms which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, straight-chain or branched alkoxy having up to 6 carbon atoms, or by a group of the formula $-NR^{51'}R^{52'}$,




in which


$R^{51'}$ and $R^{52'}$ have the meaning of R^{51} and R^{52} given above and are identical to or different from this meaning,

and/or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by aryl having 6 to 10 carbon atoms or by a 5- to 7-membered saturated, partially unsaturated or unsaturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, if appropriate also attached via an N-function, where the ring systems for their part may be substituted by hydroxyl or by straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

or

R^3 and R^4 together with the nitrogen atom form radicals of the formulae





R^5 and R^6 are identical or different and represent hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or represent straight-chain or branched alkoxy having up to 6 carbon atoms,

5 and their respective salts, hydrates, alkoxides and tautomers.

Among the compounds mentioned in WO-A-99/24433, very particular preference according to the invention is given to 2-[2-ethoxy-5-(4-ethyl-piperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one (Ex. 19 of
10 WO-A-99/24433) and its salts, such as, for example, the hydrochloride (Ex. 20 of WO-A-99/24433), in particular in the form of the trihydrate (Ex. 336 of WO-A-99/24433).

Further preferred PDE V inhibitors according to the invention are (a) zaprinast (*Am. J. Physiol.*, Vol. 264, February 1993, pages H419-H422; *The Journal of Urology*, Vol. 147, pages 1650-1655 (June 1992); *The New England Journal of Medicine*, Vol. 326(2), pages 90-94 (9 January 1992)); (b) propentofylline (JP-A-03/044324); and (c) pentoxifylline (*Postgraduate Medicine*, Vol. 93, No. 3, Impotence, 15
February 1993, pages 65-72; *J.A.G.S.*, 41, pages 363-366, 1993).

20

Further PDE V inhibitors which are preferred according to the invention are disclosed in the following publications, the contents of which are included in their entirety by way of reference:

- 5-substituted pyrazolo-[4,3-d]-pyrimidin-7-one according to EP-A-0 201 188;
- 25 • griseolic acid derivatives according to EP-A-0 214 708 and EP-A-0 319 050;
- 2-phenylpurinone derivatives according to EP-A-0 293 063;
- phenylpyridone derivatives according to EP-A-0 347 027;
- fused pyrimidine derivatives according to EP-A-0 347 146;

- condensed pyrimidine derivatives according to EP-A-0 349 239;
- pyrimidopyrimidine derivatives according to EP-A-0 351 058;
- purine compounds according to EP-A-0 352 960;
- quinazoline derivatives according to EP-A-0 371 731;
- 5 • phenylpyrimidone derivatives according to EP-A-0 395 328;
- imidazoquinoxalinone derivatives or their aza analogs according to EP-A-0 400 583;
- phenylpyrimidone derivatives according to EP-A-0 400 799;
- phenylpyridone derivatives according to EP-A-0 428 268;
- 10 • pyrimidopyrimidine derivatives according to EP-A-0 442 204;
- 4-aminoquinazoline derivatives according to EP-A-0 579 496;
- 4,5-dihydro-4-oxo-pyrrolo-[1,2-a]-quinoxaline derivatives and their aza analogs according to EP-A-0 584 487;
- polycyclic guanine derivatives according to WO-A-91/19717;
- 15 • nitrogen-containing heterocycles according to WO-A-93/07124;
- polycyclic 2-benzyl-guanine derivatives according to WO-A-94/19351;
- quinazoline derivatives according to US-A-4 060 615;
- 6-heterocycl-yl-pyrazolo-[3,4-d]-pyrimidin-4-ones according to US-A-5 294 612;
- benzimidazoles according to JP-A 5-222000;
- 20 • cycloheptimidazole according to *European Journal of Pharmacology* 1994, **251**, page 1;
- N-containing heterocycles according to WO-A-94/22855.

Further PDE inhibitors which are preferred and can be used according to the invention are:

- tetracyclic derivatives according to WO-A-95/19978;
- pyrazolopyrimidine derivatives according to EP-A-0 636 626;
- 4-aminopyrimidine derivatives according to EP-A-0 640 599;
- imidazoquinazoline derivatives according to EP-A-0 668 280;
- 30 • quinazoline compounds according to EP-A-0 669 324;
- 4-aminoquinazoline derivatives according to US-A-5 436 233.

Further PDE inhibitors which are likewise preferred and can be used according to the invention are the compounds according to EP-A-0 579 496; WO-A-93/07124; US-A-5 294 612 and WO-A-94/22855, the contents of which are hereby included by way of reference.

5

Examples of PDE inhibitors from the publications mentioned above which are particularly preferred according to the invention are the following compounds:

- 1,3-dimethyl-5-benzylpyrazolo-[4,3-d]-pyrimidin-7-one (preparation according to EP-A-0 201 188, Example 1);
- 10 • 2-(2-propoxyphenyl)-6-purinone (preparation according to EP-A-0 293 063, Example 1);
- 6-(2-propoxyphenyl)-1,2-dihydro-2-oxopyridine-3-carboxamide (preparation according to EP-A-0 347 027, Example 2);
- 15 • 2-(2-propoxyphenyl)-pyrido-[2,3-d]-pyrimid-4(3H)-one (preparation according to EP-A-0 347 146, Example 1);
- 7-methylthio-4-oxo-2-(2-propoxyphenyl)-3,4-dihydropyrimido-[4,5-d]-pyrimidine (preparation according to EP-A-0 351 058, Example 1);
- 6-hydroxy-2-(2-propoxyphenyl)-pyrimidine-4-carboxamide (preparation according to EP-A-0 395 328, Example 15);
- 20 • 1-ethyl-3-methylimidazo-[1,5-a]-quinoxalin-4(5H)-one (preparation according to EP-A-0 400 583);
- 4-phenylmethylamino-6-chloro-2-(1-imidazolyl)-quinoxaline (preparation according to EP-A-0 579 496, Example 5(c));
- 5-ethyl-8-[3-(N-cyclohexyl-N-methylcarbamyl)-propyloxy]-4,5-dihydro-4-oxo-
25 pyrido-[3,2-e]-pyrrolo-[1,2-a]-pyrazine (preparation according to EP-A-0 584 487, Example 1);
- 5'-methyl-3'-(phenylmethyl)-spiro-[cyclopentane-1,7'(8'H)-(3'H)-imidazo[2,1-b]-purin]-4'(5'H)-one (preparation according to WO-A-91/19717, Example 9A3);
- 30 • 1-[6-chloro-4-(3,4-methylenedioxybenzyl)-aminoquinazolin-2-yl]-piperidine-4-carboxylic acid (preparation according to WO-A-93/97124);
- (6aR,9aS)-2-(4-trifluoromethylphenyl)-methyl-5-methyl-3,4,5,6a,7,8,9,9a-

octahydrocyclopent-[4,5]-imidazo-[2,1-b]-purin-4-one (preparation according to WO-A-94/19351, Example 14);

- 1-tert-butyl-3-phenylmethyl-6-(4-pyridyl)-pyrazolo-[3,4-d]-pyrimid-4-one (preparation according to US-A-5 294 612, Example 90);
- 5 • 1-cyclopentyl-3-methyl-6-(4-pyridyl)-4,5-dihydro-1H-pyrazolo-[3,4-d]-pyrimid-4-one (preparation according to US-A-5 294 612, Example 83);
- 2-butyl-1-(2-chlorobenzyl)-6-ethoxycarbonylbenzimidazole (preparation according to JP-A-5-222000);
- 2-(4-carboxypiperidino)-4-(3,4-methylenedioxybenzyl)-amino-6-nitroquinazoline
- 10 (preparation according to WO-A-94/22855, Example II);
- 2-phenyl-8-ethoxycycloheptimidazole (KT2-734);
- 1-[6-chloro-4-(3,4-methyldioxybenzyl)-aminoquinazolin-2-yl]-piperidine-4-carboxylic acid (preparation according to WO-A-93/07124);
- (6aR,9aS)-2-(4-trifluoromethylphenyl)-methyl-5-methyl-3,4,5,6a,7,8,9,9a-
- 15 octahydrocyclopent-[4,5]-imidazo-[2,1-b]-purin-4-one (preparation according to WO-A-94/19351, Example 14);
- 1-tert-butyl-3-phenylmethyl-6-(4-pyridyl)-pyrazolo-[3,4-d]-pyrimid-4-one (preparation according to US-A-5 294 612, Example 90);
- 1-cyclopentyl-3-methyl-6-(4-pyridyl)-4,5-dihydro-1H-pyrazolo-[3,4-d]-pyrimid-
- 20 4-one (preparation according to US-A-5 294 612, Example 83);
- 2-(4-carboxypiperidino)-4-(3,4-methylenedioxybenzyl)-amino-6-nitro-quinazoline (preparation according to WO-A-94/22855, Example II).

Further cGMP PDE inhibitors which can be used according to the invention are the

25 following compounds:

- pyrazolopyrimidine derivatives according to EP-A-0 636 626;
- 4-aminopyrimidine derivatives according to EP-A-0 640 599;
- imidazoquinazoline derivatives according to WO-A-95/06648;
- anthranilic acid derivatives according to WO-A-95/18097;
- 30 • 4-aminoquinazoline derivatives according to US-A-5 436 233;
- tetracyclic derivatives according to WO-A-95/19978;
- imidazoquinazoline derivatives according to EP-A-0 668 280;

- quinazoline compounds according to EP-A-0 669 324.

Further PDE inhibitors which are preferred according to the invention, in particular cGMP PDE inhibitors, are disclosed in the following publications, whose contents are hereby included by way of reference:

- quinolone derivatives according to WO-A-98/53819;
- pyrazolo[4,3-d]-pyrimidine derivatives according to EP-A-911333;
- pyrazolopyrimidones according to WO-A-98/49166;
- compounds according to WO-A-96/16644;
- 10 • bicyclic heterocycles according to WO-A-96/16657;
- pyrazolopyrimidones according to WO-A-93/06104;
- pyrazolopyrimidones according to WO-A-93/07149;
- quinazolines according to WO-A-93/12093;
- purinones according to WO-A-94/00453;
- 15 • pyridopyrimidinones according to WO-A-94/05661;
- compounds according to US-A-5 272 147 and US-A-5 250 534; and
- quinazolinones according to WO-A-93/12095.

Further PDE inhibitors which are preferred according to the invention, in particular cGMP PDE inhibitors, are selected from the group consisting of:

- 4-(3-chloro-4-methoxybenzyl)amino-1-(4-hydroxypiperidino)-6-phthalazine-carbonitrile and its salts, tautomers, alkoxides and hydrates, in particular 4-(3-chloro-4-methoxybenzyl)-amino-1-(4-hydroxypiperidino)-6-phthalazine-carbonitrile hydrochloride;
- 25 • 1-[4-[(1,3-benzodioxol-5-ylmethyl)amino]-6-chloro-2-quinazolyl]-4-piperidine-carboxylic acid and its salts, tautomers, alkoxides and hydrates, in particular the sodium salt;
- 4-bromo-6-[3-(4-chlorophenyl)propoxy]-5-[(3-piperidinylmethyl)amino]-3(2H)-pyridazinone and its salts, tautomers, alkoxides and hydrates, in particular
- 30 4-bromo-6-[3-(4-chlorophenyl)propoxy]-5-[(3-piperidinylmethyl)-amino]-3(2H)-pyridazinone hydrochloride.

Further PDE inhibitors which are preferred according to the invention, in particular cGMP PDE inhibitors, are disclosed in the following publications, whose contents are hereby included by way of reference:

- carbazole derivatives and their salts according to WO-A-99/21831 (Fujisawa);
- 5 • dihydropyrazolopyrimidinone derivatives according to WO-A-98/40384 (BAYER) and DE-A-197 09 877 (BAYER);
- purine and pyrazolopyrimidine derivatives according to DE-A-197 09 126 (BAYER);
- isoquinolinone derivatives according to WO-A-98/38168 (Tanabe);
- 10 • substituted azaindole compounds according to JP-A-10120681 (Fujisawa);
- indolizine compounds according to JP-A-10120680 (Fujisawa);
- (amino-thieno-pyrimidinyl)-heterocyclic acid derivatives according to DE-A-196 44 228 (Merck) and WO-A-98/17668 (Merck);
- 2-amino-thiophene-3-carboxamide derivatives according to DE-A-196 42 451 (Merck) and WO-A-98/16521 (Merck);
- 15 • pyridocarbazole derivatives according to WO-A-97/45427 (Mochida);
- purinone derivatives according to EP-A-771799 (BAYER);
- N-benzyl-3-indenylacetamide derivatives according to WO-A-99/31065 (Baverstock/Cell Pathways/Univ. Arizona State);
- 20 • pyridocarbazole compounds according to WO-A-99/26946 (Mochida);
- thienopyrimidine derivatives according to DE-A-197 52 952 (Merck) and WO-A-99/28325 (Merck);
- indole derivatives according to WO-A-98/15530 (Fujisawa);
- pyrazolopyridazinones according to WO-A-98/14448 (Kyorin);
- 25 • phenylalkylthienopyrimidine derivatives according to DE-A-196 32 423 (Merck) and WO-A-98/06722 (Merck);
- compounds according to WO-A-96/21435 (Euroceltique);
- dihydropyrazolopyrroles according to WO-A-95/19362 (Pfizer);
- compounds according to WO-A-95/00516 (Euroceltique).

30

In addition to the two active compound components A and B mentioned above, the combination preparation according to the invention may comprise any further active

compounds, provided these compounds are not contraindicated for this area of indication and do not adversely affect the action of the PDE inhibitor and the antilipemic.

- 5 These further optionally present active compounds, like the active compound components A and B, can be present either as a true mixture together with A and/or B or else spatially separated therefrom. They can be administered concomitantly or simultaneously or successively with the active compound component(s) A and/or B.
- 10 The further optionally present active compounds of the combination preparation according to the invention include, for example:
- further active compounds which improve the capability to maintain an erection and which do not belong to the class of the PDE inhibitors, such as, for example: α -adrenergic antagonists, such as, for example, yohimbine or Vasomax[®] from
 - 15 Zonagen; or else substances as mentioned in WO-A-98/52569, the content of which is hereby included by way of reference; or prostaglandine-E1; or serotonin antagonists;
 - active compounds from the cardiovascular field of indication;
 - active compounds from the CNS and cerebral field of indication;
 - 20 • vitamins;
 - minerals;
 - trace elements.

Suitable administration forms for the administration of the two active compound

25 components A and B (and any other active compounds present) are in each case all customary administration forms. The administration is preferably carried out orally, perlingually, sublingually, nasally, transdermally, buccally, intravenously, rectally, inhalatively or parenterally. The administration is preferably carried out orally, sublingually or nasally. Very particular preference is given to oral administration.

30

In the case of spatially separated and/or successive administration, the two active compound components A and B can also be administered in different administration

forms.

The two active compound components A and B – together or spatially separated – can in each case be converted in a manner known per se into customary formulations
5 such as tablets including coated tablets, pills, granules, aerosols, syrups, emulsions, suspensions and solutions, using inert, non-toxic, pharmaceutically suitable carriers or solvents. In this case, the therapeutically active components A and B should in each case be present in a concentration from approximately 0.5 to 90% by weight of the total mixture, i.e. in amounts which are sufficient in order to achieve the dosage
10 range indicated.

The formulations are prepared, for example, by extending the two active compound components A and B with solvents and/or carriers, if appropriate using emulsifying agents and/or dispersing agents, it being possible, for example if the diluent used is
15 water, to use organic solvents as auxiliary solvents, if appropriate.

For use in humans, in the case of oral administration, dosages of from 0.001 to 50 mg/kg, preferably from 0.001 mg/kg to 20 mg/kg, in particular from 0.001 to 10 mg/kg, of body weight, particularly preferably from 0.001 mg/kg to 5 mg/kg, of
20 the respective active compound component A or B are administered to obtain effective and useful results.

In spite of this, it may be necessary, if appropriate, to depart from the amounts mentioned, namely depending on the body weight and/or on the type of
25 administration route, on the individual response toward the combination preparation, the manner of its formulation and the time or interval at which administration takes place. Thus, in some cases it may be adequate to manage with less than the abovementioned minimum amount, while in other cases the upper limit mentioned must be exceeded.

30

In the case of the administration of relatively large amounts, it may be advisable to divide these into several individual administrations over the course of the day.

The present invention is illustrated by the embodiment below in a purely exemplary manner, and by no means limited thereby.

Example

In a placebo-controlled study, twelve adult male rabbits of the animal strain HsdHHL Watanables having a body weight of 3-5 kg, which, owing to breeding, suffer from hypercholesterolemia and arteriosclerosis, were treated with a selective cGMP PDE inhibitor (3 mg/kg, intravenously).

The animals had free access to water and, for 2 hours per day, to feed. The animals were kept in a 10/14 hour day/night rhythm (light on from 8 am onwards), and the ambient temperature was 22 to 24°C.

The selective cGMP PDE inhibitor used was 2-[2-ethoxy-5-(4-ethyl-piperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one hydrochloride trihydrate (Ex. 336 of WO-A-99/24433).

For intravenous injection, the PDE inhibitor was dissolved in physiological saline (0.9% by volume).

For seven days prior to the treatment with the PDE inhibitor, i.e. for 7 successive days, an antilipemic, i.e. cerivastatin sodium (i.e. the monosodium salt of cerivastatin) was administered subcutaneously to six of the twelve rabbits (1 mg/kg of body weight, cerivastatin sodium dissolved in physiological saline).

Instead of this, a physiological saline was administered subcutaneously to the other six rabbits (control).

In each case 1 day and 3 days after the 7-day pretreatment with cerivastatin sodium had ended, the six rabbits which had been pretreated with cerivastatin sodium were given the PDE inhibitor.

The other six rabbits, which had not been pretreated with cerivastatin sodium, were given the PDE inhibitor 1 day after the 7-day subcutaneous administration of the saline (control) had ended.

The erection was evaluated by measuring the length of the protruding penis using a vernier caliper. The measurement was carried out in each case 5, 10, 15, 30, 45, 60 and 120 minutes after the administration of the PDE inhibitor. To this end, the animals were in each case removed from the cage, held by neck hair and hind legs, turned onto their back and measured. Under rest conditions, i.e. in the non-erect state, the penis of the rabbit is not visible in the pubic region and completely covered by the penis skin.

10 As illustrated by the attached **Fig. 2**, the six rabbits which had been pretreated with the antilipemic cerivastatin sodium (upper and middle curves) showed, after administration of the PDE inhibitor, surprisingly a considerably more pronounced erection than the six rabbits which had not been pretreated with the antilipemic (lower curve).

15 Comparison of the upper and middle curves of **Fig. 2** also shows that the enhanced capability to sustain an erection in the pretreated animals decreased again once the treatment with the antilipemic was stopped. Thus, the erection 1 day after the cerivastatin sodium pretreatment had ended was still more pronounced than even 3 days after the cerivastatin sodium pretreatment had ended. This too confirms that the surprising improvement in the activity of the PDE inhibitor is caused by the antilipemic.

25 As shown by the experiment, even a pretreatment with the antilipemic which is initiated shortly before the administration of the PDE inhibitor causes, entirely unexpectedly, an enhanced PDE-inhibitor-triggered erection. In this manner, the activity of the PDE inhibitor is surprisingly improved, since, in view of the short pretreatment interval with the antilipemic, no (significant) improvement in the hypercholesterolemia, not to mention the arteriosclerosis, had occurred.

30 Without wishing to adhere to a certain theory, the synergistic effect of the PDE inhibitor together with the antilipemic can possibly be explained by an improved endothelium function, as already discussed above in the general description. This

synergistic effect, however, was totally unforeseeable for the person skilled in the art and therefore has to be considered to be entirely surprising.

Thus, the embodiment confirms in an impressive manner the improved PDE-
5 inhibitory action of the PDE inhibitor by the antilipemic administered in combination therewith.


However, the person skilled in the art will be familiar with numerous further
10 embodiments when practicing the present invention, without being outside of the scope of the present invention.

Patent claims

1. A combination preparation, comprising as pharmaceutically active ingredients at least one active compound component A and at least one active compound component B, characterized in that the active compound component A is a PDE inhibitor, preferably a cGMP PDE inhibitor, and the active compound component B is an antilipemic.
2. The combination preparation as claimed in claim 1 for treating sexual dysfunction in men or women.
3. The combination preparation as claimed in claim 2 for treating erectile dysfunction.
4. The combination preparation as claimed in any of claims 1 to 3, characterized in that the two active compound components A and B are used either simultaneously or successively.
5. The combination preparation as claimed in any of claims 1 to 4, characterized in that the active compound components A and B are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
6. The combination preparation as claimed in any of claims 1 to 4, characterized in that the active compound components A and B are (spatially) separated, in particular as a kit-of-parts.
7. The combination preparation as claimed in any of claims 1 to 6, characterized in that the antilipemic (active compound component B) is selected from the group consisting of (a) HMG-CoA-reductase inhibitors; (b) squalene synthase inhibitors; (c) bile acid sequestrants; (d) fibric acid and its derivatives; (e) nicotinic acid and its analogs; (f) ω 3-fatty acids.
8. The combination preparation as claimed in claim 7, characterized in that the

antilipemic (active compound component B) is an HMG-CoA-reductase inhibitor and is in particular selected from the group of the statins, preferably from the group consisting of atorvastatin, cerivastatin, fluvastatin, lovastatin, pravastatin, itavastatin, simvastatin and (+)-(3R,5S)-bis-(7-(4-(4-fluorophenyl)-6-isopropyl-2-(N-methyl-N-methanesulfonylamino)-pyrimidin-5-yl)-3,5-dihydroxy-6(E)-heptenoic acid, and their respective salts, hydrates, alkoxides, esters and tautomers.

9. The combination preparation as claimed in claim 8, characterized in that the antilipemic (active compound component B) is atorvastatin or its salt, hydrate, alkoxide, ester and tautomer.
10. The combination preparation as claimed in claim 8, characterized in that the antilipemic (active compound component B) is cerivastatin or its salt, hydrate, alkoxide, ester and tautomer.
11. The combination preparation as claimed in any of claims 1 to 10, characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and is in particular selected from the group consisting of pyrazolopyrimidones of the general formula below




in which

- R^1 represents hydrogen; C_1 - C_3 -alkyl; C_1 - C_3 -perfluoroalkyl; or C_3 - C_5 -cycloalkyl;
- R^2 denotes hydrogen; C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl; C_1 - C_3 -perfluoroalkyl; or C_3 - C_6 -cycloalkyl;
- R^3 is C_1 - C_6 -alkyl, optionally substituted by C_3 - C_6 -cycloalkyl; C_1 - C_6 -perfluoroalkyl, C_3 - C_5 -cycloalkyl; C_3 - C_6 -alkenyl; or C_3 - C_6 -alkinyl;

- 5 R^4 represents C_1 - C_4 -alkyl, optionally substituted by OH, NR^5R^6 , CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkenyl, optionally substituted by CN, $CONR^5R^6$ or CO_2R^7 ; C_2 - C_4 -alkanoyl, optionally substituted by NR^5R^6 ; (hydroxy)- C_2 - C_4 -alkyl, optionally substituted by NR^5R^6 , (C₂-C₃-alkoxy)- C_1 - C_2 -alkyl, optionally substituted by OH or NR^5R^6 , CO_2R^7 ; halogen; NR^5R^6 , $NHSO_2NR^5R^6$; $NHSO_2R^8$; $SO_2NR^9R^{10}$; or phenyl, pyridyl, pyrimidinyl, imidazolyl, oxazolyl, thiazolyl, thienyl or triazolyl, each of which is optionally substituted by methyl;
- 10 R^5 and R^6 each independently of one another denote hydrogen or C_1 - C_4 -alkyl; or together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4-N(R^{11})-piperazinyl or imidazolyl group, where this group is optionally substituted by methyl or OH;
- 15 R^7 is hydrogen or C_1 - C_4 -alkyl;
- R^8 represents C_1 - C_3 -alkyl, optionally substituted by NR^5R^6 ;
- 20 R^9 and R^{10} together with the nitrogen atom to which they are attached form a pyrrolidinyl, piperidino, morpholino, 4-N(R^{12})-piperazinyl group, where this group is optionally substituted by C_1 - C_4 -alkyl, C_1 - C_3 -alkoxy, $NR^{13}R^{14}$ or $CONR^{13}R^{14}$;
- 25 R^{11} denotes hydrogen, C_1 - C_3 -alkyl, optionally substituted by phenyl; (hydroxy)- C_2 - C_3 -alkyl; or C_1 - C_4 -alkanoyl;
- R^{12} is hydrogen, C_1 - C_6 -alkyl, (C₁-C₃-alkoxy)- C_2 - C_6 -alkyl; (hydroxy)- C_2 - C_6 -alkyl; ($R^{13}R^{14}N$)- C_2 - C_6 -alkyl; ($R^{13}R^{14}NOC$)- C_1 - C_6 -alkyl; $CONR^{13}R^{14}$; $CSNR^{13}R^{14}$, or $C(NH)NR^{13}R^{14}$; and
- 30 R^{13} and R^{14} each independently of one another represent hydrogen; C_1 - C_4 -alkyl; (C₁-C₃-alkoxy)- C_2 - C_4 -alkyl; or (hydroxy)- C_2 - C_4 -alkyl;

and their respective salts, hydrates, alkoxides and tautomers.

12. The combination preparation as claimed in any of claims 1 to 10,
 5 characterized in that the PDE inhibitor (active compound component A) is a
 cGMP PDE inhibitor and is in particular selected from the group consisting
 of 2-phenyl-substituted imidazotriazinones of the general formula



in which

10

R^1 represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms;

R^2 represents straight-chain alkyl having up to 4 carbon atoms;


15

R^3 and R^4 are identical or different and represent hydrogen or represent straight-chain or branched alkenyl or alkoxy having in each case up to 8 carbon atoms, or

20

represent a straight-chain or branched alkyl chain having up to 10 carbon atoms which is optionally interrupted by an oxygen atom and which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of trifluoromethyl, trifluoromethoxy, hydroxyl, halogen, carboxyl, benzyloxycarbonyl, straight-chain or branched alkoxycarbonyl having up to 6 carbon atoms and/or by radicals of the formulae $-SO_3H$, $-(A)_a-NR^7R^8$, $-O-CO-NR^7R^8$, $-S(O)_b-R^9$, $-P(O)(OR^{10})(OR^{11})$,

25



in which

a and b are identical or different and represent a number 0 or 1,

5

A represents a radical CO or SO₂,

R⁷, R^{7'}, R⁸ and R^{8'} are identical or different and represent hydrogen, or
 represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10
 carbon atoms, a 5- to 6-membered unsaturated, partially unsaturated
 or saturated optionally benzo-fused heterocycle having up to 3
 heteroatoms from the group consisting of S, N and O, where the
 abovementioned ring systems are optionally mono- to polysubstituted
 by identical or different substituents from the group consisting of
 hydroxyl, nitro, trifluoromethyl, trifluoromethoxy, carboxyl, halogen,
 straight-chain or branched alkoxy or alkoxycarbonyl having in each
 case up to 6 carbon atoms or by a group of the formula
 -(SO₂)_c-NR¹²R¹³,

10

15

20

in which

c represents a number 0 or 1,

R^{12} and R^{13} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 5 carbon atoms,

5 or

R^7 , $R^{7'}$, R^8 and $R^{8'}$ represent straight-chain or branched alkoxy having up to 6 carbon atoms, or

represent straight-chain or branched alkyl having up to 8 carbon atoms which is optionally mono- or polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, aryl having 6 to 10 carbon atoms, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms, or by a group of the formula $-(CO)_d-NR^{14}R^{15}$,

15 in which

R^{14} and R^{15} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

20 and

d represents a number 0 or 1,

25 or

R^7 and R^8 and/or $R^{7'}$ and $R^{8'}$ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O or a radical of the formula $-NR^{16}$,

30 in which

R¹⁶ represents hydrogen, aryl having 6 to 10 carbon atoms, benzyl, a 5- to 7-membered aromatic or saturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, which heterocycle is optionally substituted by methyl, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl,

R⁹ represents aryl having 6 to 10 carbon atoms, or represents straight-chain or branched alkyl having up to 4 carbon atoms,

R¹⁰ and R¹¹ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

and/or the alkyl chain listed above under R³/R⁴ is optionally substituted by cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or by a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula -NR¹⁷,

in which

R¹⁷ represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain or branched acyl or alkoxy having in each case up to 4 carbon atoms, or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl and straight-chain or branched alkoxy having up to 6 carbon atoms,

and where aryl and the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of nitro, halogen, -SO₃H, straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms, hydroxyl, trifluoromethyl, trifluoromethoxy and/or by a radical

of the formula $-\text{SO}_2\text{NR}^{18}\text{R}^{19}$,

in which

5 R^{18} and R^{19} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms,

and/or



10 R^3 or R^4 represent a group of the formula $-\text{NR}^{20}\text{R}^{21}$,

in which

15 R^{20} and R^{21} have the meaning of R^{18} and R^{19} given above and are identical to or different from this meaning,

and/or

20 R^3 or R^4 represent adamantyl, or represent radicals of the formulae



25 or represent cycloalkyl having 3 to 8 carbon atoms, aryl having 6 to 10 carbon atoms or represent a 5- to 7-membered partially unsaturated, saturated or unsaturated optionally benzo-fused

heterocycle which may contain up to 4 heteroatoms from the group consisting of S, N; O or a radical of the formula $-NR^{22}$,

in which


R^{22} has the meaning of R^{16} given above and is identical to or different from this meaning, or represents carboxyl, formyl or straight-chain or branched acyl having up to 5 carbon atoms,

and where cycloalkyl, aryl and/or the heterocycle are optionally mono- to polysubstituted by identical or different substituents from the group consisting of halogen, triazolyl, trifluoromethyl, trifluoromethoxy, carboxyl, straight-chain or branched acyl or alkoxy carbonyl having in each case up to 6 carbon atoms, nitro, and/or by groups of the formulae $-SO_3H$, $-OR^{23}$, $(SO_2)_eNR^{24}R^{25}$, $-P(O)(OR^{26})(OR^{27})$,

in which

e represents a number 0 or 1,

R^{23} represents a radical of the formula



represents cycloalkyl having 3 to 7 carbon atoms, or represents hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms which is optionally substituted by cycloalkyl having 3 to 7 carbon atoms, benzyloxy, tetrahydropyranyl, tetrahydrofuranyl, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms, carboxyl, benzyloxycarbonyl or phenyl which for its part may be mono- to polysubstituted by identical or

different substituents from the group consisting of straight-chain or branched alkoxy having up to 4 carbon atoms, hydroxyl and halogen,

and/or alkyl is optionally substituted by radicals of the formulae -CO-NR²⁸R²⁹ or -CO-R³⁰,

in which

R²⁸ and R²⁹ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 8 carbon atoms, or

R²⁸ and R²⁹ together with the nitrogen atom form a 5- to 7-membered saturated heterocycle which may optionally contain a further heteroatom from the group consisting of S and O,

and

R³⁰ represents phenyl or adamantyl,

R²⁴ and R²⁵ have the meaning of R¹⁸ and R¹⁹ given above and are identical to or different from this meaning,

R²⁶ and R²⁷ have the meaning of R¹⁰ and R¹¹ given above and are identical to or different from this meaning

and/or cycloalkyl, aryl and/or the heterocycle are optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl, carboxyl, by a 5- to 7-membered heterocycle having up to 3 heteroatoms from the group consisting of S, N and O or by groups of the formula -SO₂-R³¹, P(O)(OR³²)(OR³³) or -NR³⁴R³⁵,

in which

R^{31} is hydrogen or has the meaning of R^9 given above and is identical to or different from this meaning,

5 R^{32} and R^{33} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{34} and R^{35} are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 6 carbon atoms which is optionally substituted by hydroxyl or straight-chain or branched alkoxy having
10 up to 4 carbon atoms, or

R^{34} and R^{35} together with the nitrogen atom form a 5- to 6-membered saturated heterocycle which may contain a further heteroatom from
15 the group consisting of S and O or a radical of the formula $-NR^{36}$,

in which

R^{36} represents hydrogen, hydroxyl, straight-chain or branched
20 alkoxy carbonyl having up to 7 carbon atoms or straight-chain or branched alkyl having up to 5 carbon atoms which is optionally substituted by hydroxyl,

or

25 R^3 and R^4 together with the nitrogen atom form a 5- to 7-membered unsaturated or saturated or partially unsaturated optionally benzo-fused heterocycle which may optionally contain up to 3 heteroatoms from the group consisting of S, N, O or a radical of the formula $-NR^{37}$,

30

in which

R^{37} represents hydrogen, hydroxyl, formyl, trifluoromethyl, straight-chain

or branched acyl, alkoxy or alkoxycarbonyl having in each case up to 4 carbon atoms,

or represents straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, trifluoromethyl, carboxyl, straight-chain or branched alkoxy or alkoxycarbonyl having in each case up to 6 carbon atoms or by groups of the formula - (D)_f-NR³⁸R³⁹, -CO-(CH₂)_g-O-CO-R⁴⁰, -CO-(CH₂)_h-OR⁴¹ or -P(O)(OR⁴²)(OR⁴³),

in which

g and h are identical or different and represent a number 1, 2, 3 or 4,

and

f represents a number 0 or 1,

D represents a group of the formula -CO or -SO₂,

R³⁸ and R³⁹ are identical or different and have the meaning of R⁷ and R⁸ given above,

R⁴⁰ represents straight-chain or branched alkyl having up to 6 carbon atoms,

R⁴¹ represents straight-chain or branched alkyl having up to 6 carbon atoms,

R⁴² and R⁴³ are identical or different and represent hydrogen or straight-chain or branched alkyl having up to 4 carbon atoms,

or

R^{37} represents a radical of the formula $-(CO)_i-E$,

in which

i represents a number 0 or 1,


E represents cycloalkyl having 3 to 7 carbon atoms or benzyl,
represents aryl having 6 to 10 carbon atoms or a 5- to 6-
membered aromatic heterocycle having up to 4 heteroatoms
from the group consisting of S, N and O, where the ring
systems listed above are optionally mono- to polysubstituted
by identical or different substituents from the group consisting
of nitro, halogen, $-SO_3H$, straight-chain or branched alkoxy
having up to 6 carbon atoms, hydroxyl, trifluoromethyl,
trifluoromethoxy or by a radical of the formula $-SO_2-NR^{44}R^{45}$,

in which

R^{44} and R^{45} have the meaning of R^{18} and R^{19} given above and
are identical to or different from this meaning,


or

E represents radicals of the formulae



and the heterocycle listed under R^3 and R^4 , which is formed together

with the nitrogen atom, is optionally mono- to polysubstituted by identical or different substituents, if appropriate also geminally, by hydroxyl, formyl, carboxyl, straight-chain or branched acyl or alkoxy carbonyl having in each case up to 6 carbon atoms, nitro and groups of the formulae $-P(O)(OR^{46})(OR^{47})$,



$=NR^{48}$ or $-(CO)_jNR^{49}R^{50}$,

in which

R^{46} and R^{47} have the meaning of R^{10} and R^{11} given above and are identical to or different from this meaning,

R^{48} is hydroxyl or straight-chain or branched alkoxy having up to 4 carbon atoms,

j is a number 0 or 1,

and

R^{49} and R^{50} are identical or different and have the meaning of R^{14} and R^{15} given above,

and/or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by straight-chain or branched alkyl having up to 6 carbon atoms which is optionally mono- to polysubstituted by identical or different substituents from the group consisting of hydroxyl, halogen, carboxyl, cycloalkyl or cycloalkyloxy having in each case 3 to 8 carbon atoms, straight-chain or branched alkoxy or alkoxy carbonyl having in each case up to 6 carbon atoms or by a radical of the formula $-SO_3H$, $-NR^{51}R^{52}$ or $P(O)OR^{53}OR^{54}$,

in which

5 R^{51} and R^{52} are identical or different and represent hydrogen, phenyl, carboxyl, benzyl or straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

10 R^{53} and R^{54} are identical or different and have the meaning of R^{10} and R^{11} given above,

and/or the alkyl is optionally substituted by aryl having 6 to 10 carbon atoms which for its part may be mono- to polysubstituted by identical or different substituents from the group consisting of halogen, hydroxyl, straight-chain or branched alkoxy having up to 6 carbon atoms, or by a group of the formula $-NR^{51'}R^{52'}$,


in which

20 $R^{51'}$ and $R^{52'}$ have the meaning of R^{51} and R^{52} given above and are identical to or different from this meaning,

25 and/or the heterocycle listed under R^3 and R^4 , which is formed together with the nitrogen atom, is optionally substituted by aryl having 6 to 10 carbon atoms or by a 5- to 7-membered saturated, partially unsaturated or unsaturated heterocycle having up to 3 heteroatoms from the group consisting of S, N and O, if appropriate also attached via an N-function, where the ring systems for their part may be substituted by hydroxyl or by straight-chain or branched alkyl or alkoxy having in each case up to 6 carbon atoms,

30 or

R^3 and R^4 together with the nitrogen atom form radicals of the formulae



R^5 and R^6 are identical or different and represent hydrogen, straight-chain or branched alkyl having up to 6 carbon atoms, hydroxyl or represent straight-chain or branched alkoxy having up to 6 carbon atoms.

- 5 and their respective salts, hydrates, alkoxides and tautomers.
13. The combination preparation as claimed in any of claims 1 to 10, characterized in that the PDE inhibitor (active compound component A) is a cGMP PDE inhibitor and in particular is selected from the group consisting of (a) 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one (sildenafil) and its salts, hydrates, alkoxides and tautomers; and (b) 2-[2-ethoxy-5-(4-ethylpiperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f]-[1,2,4]-triazin-4-one and its salts, hydrates, alkoxides and tautomers.
- 15
14. The combination preparation as claimed in claim 13, in that the PDE inhibitor (active compound component A) is 5-[2-ethoxy-5-(4-methyl-1-piperazinylsulfonyl)-phenyl]-1-methyl-3-n-propyl-1,6-dihydro-7H-pyrazolo-[4,3-d]-pyrimidin-7-one citrate (sildenafil citrate, ViagraTM) or 2-[2-ethoxy-5-(4-ethylpiperazine-1-sulfonyl)-phenyl]-5-methyl-7-propyl-3H-imidazo[5,1-f][1,2,4]triazin-4-one hydrochloride trihydrate.
- 20
15. The use of antilipemics for enhancing the activity of PDE inhibitors, in

particular cGMP PDE inhibitors.

- 5
16. The use as claimed in claim 15 in the treatment of sexual dysfunction in men and women, in particular in the treatment of erectile dysfunction.
17. The use as claimed in claim 15 or 16, characterized in that the antilipemic and the PDE inhibitor are used either simultaneously or else successively.
- 10 18. The use as claimed in any of claims 15 to 17, characterized in that the antilipemic and the PDE inhibitor are present as a functional unit, in particular in the form of a mixture, a mix or a blend.
- 15 19. The use as claimed in any of claims 15 to 17, characterized in that the antilipemic and the PDE inhibitor are present (spatially) separated, in particular as a kit-of-parts.
20. The use as claimed in any of claims 15 to 19, characterized in that the antilipemic is selected from the compounds defined in claims 7 to 10.
- 20 21. The use as claimed in any of claims 15 to 20, characterized in that the PDE inhibitor is selected from the compounds defined in claims 11 to 14.

(12) NACH DEM VERTRAG ÜBER DIE INTERNATIONALE ZUSAMMENARBEIT AUF DEM GEBIET DES
PATENTWESENS (PCT) VERÖFFENTLICHTE INTERNATIONALE ANMELDUNG

(19) Weltorganisation für geistiges Eigentum
Internationales Büro



(43) Internationales Veröffentlichungsdatum
22. März 2001 (22.03.2001)

PCT

(10) Internationale Veröffentlichungsnummer
WO 01/19357 A2

(51) Internationale Patentklassifikation⁷: A61K 31/00

(74) Gemeinsamer Vertreter: BAYER AKTIENGESELLSCHAFT; 51368 Leverkusen (DE).

(21) Internationales Aktenzeichen: PCT/EP00/08836

(22) Internationales Anmeldedatum:
11. September 2000 (11.09.2000)

(25) Einreichungssprache: Deutsch

(26) Veröffentlichungssprache: Deutsch

(30) Angaben zur Priorität:
199 44 161.8 ✓ 15. September 1999 (15.09.1999) DE

(71) Anmelder (für alle Bestimmungsstaaten mit Ausnahme von US): BAYER AKTIENGESELLSCHAFT [DE/DE]; 51368 Leverkusen (DE).

(81) Bestimmungsstaaten (national): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CR, CU, CZ, DE, DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Bestimmungsstaaten (regional): ARIPO-Patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), eurasisches Patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), europäisches Patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI-Patent (BF, BJ, CF, CG, CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Veröffentlicht:

— Ohne internationalen Recherchenbericht und erneut zu veröffentlichen nach Erhalt des Berichts

Zur Erklärung der Zweibuchstaben-Codes, und der anderen Abkürzungen wird auf die Erklärungen ("Guidance Notes on Codes and Abbreviations") am Anfang jeder regulären Ausgabe der PCT-Gazette verwiesen

(72) Erfinder; und

(75) Erfinder/Anmelder (nur für US): BISCHOFF, Erwin [DE/DE]; Pahlkestr. 73, 42115 Wuppertal (DE). BISCHOFF, Hilmar [DE/DE]; Am Rohm 78, 42113 Wuppertal (DE). GIULIANO, François [FR/FR]; Centre Hospitalier, Université de Bicêtre, 78, rue de Général Leclerc, F-94275 Le Kremlin-Bicêtre (FR).

(54) Title: NOVEL COMBINATION FOR THE TREATMENT OF SEXUAL DYSFUNCTION

(54) Bezeichnung: NEUE KOMBINATION ZUR BEHANDLUNG VON SEXUELLER DYSFUNKTION

(57) Abstract: Disclosed is a combination preparation for the treatment of sexual dysfunction in men or women containing at least one active ingredient A and one active ingredient B as pharmaceutically active ingredients, whereby the active ingredient A is a PDE inhibitor, preferably a cGMP PDE inhibitor and the active ingredient B a lipid-reducing agent. Both the active ingredients A and B can be administered simultaneously or at alternate intervals, that is, as a functional unit or separated from each other.

(57) Zusammenfassung: Es wird ein Kombinationspräparat zur Behandlung der sexuellen Dysfunktion bei Männern oder Frauen beschrieben, das als pharmazeutisch wirksame Bestandteile mindestens eine Wirkstoffkomponente A und mindestens eine Wirkstoffkomponente B enthält, wobei die Wirkstoffkomponente A ein PDE-Inhibitor, vorzugsweise ein cGMP PDE-Inhibitor, und die Wirkstoffkomponente B ein Lipidsenker ist. Die beiden Wirkstoffkomponenten A und B können entweder gleichzeitig oder aber zeitlich abgestuft angewandt werden, d.h. funktionelle Einheit oder getrennt voneinander vorliegen.

WO 01/19357 A2

Fig. 1

Mechanism of action of PDE V inhibitors

Fig. 1





Fig. 2

Synergistic enhancement of the cGMP PDE-inhibitory action

Fig. 2

COMBINED DECLARATION AND POWER OF ATTORNEY

ATTORNEY DOCKET NO

Le A 33 965

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name. I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought

on the invention entitled

NOVEL COMBINATION FOR THE TREATMENT OF SEXUAL DYSFUNCTION

the specification of which is attached hereto,

or was filed on **September 11, 2000**

as a PCT Application Serial No. **PCT/EP00/08836**

I hereby state that I have reviewed and understand the contents of the above-identified specification, including the claims.

I acknowledge the duty to disclose information which is material to the patentability of this application in accordance with Title 37, Code of Federal Regulations, §1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, §119 of any foreign application(s) for patent or inventor's certificate listed below and have also identified below any foreign application for patent or inventor's certificate having a filing date before that of the application on which priority is claimed:

Prior Foreign Application(s), the priority(ies) of which is/are to be claimed:

199 44 161.8
(Number)

Germany
(Country)

September 15, 1999
(Month/Day/Year Filed)

I hereby claim the benefit under Title 35, United States Code, §120 of any United States application(s) listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States application in the manner provided by the first paragraph of Title 35, United States Code, §112, I acknowledge the duty to disclose the material information as defined in Title 37, Code of Federal Regulations, §1.56 which occurred between the filing date of the prior application and the national or PCT international filing date of this application:

(Application Serial No.)

(Filing Date)

(Status)

(patented, pending, abandoned)

(Application Serial No.)

(Filing Date)

(Status)

(patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

Le A 33 965-US

DLX

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and to transact all business in the Patent and Trademark Office connected therewith:

Jeffrey M. Greenman, Reg.No. 26,552
Barbara A. Shimei, Reg. No. 29,862
William F. Gray, Reg. No. 31,018
Alice A. Brewer, Reg. No. 32,888
Jerrie L. Chiu, Reg. No. 41,670
Susan M. Pellegrino, Reg. No. 48,972



27941

PATENT TRADEMARK OFFICE

all of Bayer Corporation, 400 Morgan Lane, West Haven, Connecticut 06516

Send Correspondence To: Mr. Jeffrey M. Greenman Bayer Corporation 400 Morgan Lane West Haven, Connecticut 06516		Direct Telephone Calls To: (203)812-3964(Jerrie L. Chiu)		
FULL NAME OF SOLE OR FIRST INVENTOR Erwin Bischoff		INVENTOR'S SIGNATURE <i>Erwin Bischoff</i>		DATE 27.02.02
RESIDENCE D 42115 Wuppertal, Germany <i>Dex</i>		CITIZENSHIP German		
POST OFFICE ADDRESS c/o BAYER AKTIENGESELLSCHAFT, D 51368 Leverkusen, Germany				
FULL NAME OF SECOND INVENTOR Hilmar Bischoff <i>Hilmar Bischoff</i>		INVENTOR'S SIGNATURE <i>Hilmar Bischoff</i>		DATE 27.02.02
RESIDENCE D 42113 Wuppertal, Germany <i>Dex</i>		CITIZENSHIP German		
POST OFFICE ADDRESS c/o BAYER AKTIENGESELLSCHAFT, D 51368 Leverkusen, Germany				
FULL NAME OF THIRD INVENTOR Francois Giuliano <i>Francois Giuliano</i>		INVENTOR'S SIGNATURE <i>Francois Giuliano</i>		DATE 16.04.02
RESIDENCE F 92210 Saint Cloud, France <i>PRX</i>		CITIZENSHIP French		
POST OFFICE ADDRESS 28, Rue Pasteur, F 92210 Saint Cloud, France				
FULL NAME OF FOURTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE		CITIZENSHIP		
POST OFFICE ADDRESS				
FULL NAME OF FIFTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE		CITIZENSHIP		
POST OFFICE ADDRESS				
FULL NAME OF SIXTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE		CITIZENSHIP		
POST OFFICE ADDRESS				
FULL NAME OF SEVENTH INVENTOR		INVENTOR'S SIGNATURE		DATE
RESIDENCE		CITIZENSHIP		
POST OFFICE ADDRESS				

Le A 33 965-US

202